

CVS Pharmacy, Inc.

-PE Product Labels –

-continued



Compare to the active ingredients in
Tylenol® Sinus Severe*

Daytime

Non-Drowsy

Severe Sinus Pain & Congestion

ACETAMINOPHEN
Pain reliever, Fever reducer

GUAIFENESIN

Expectorant

PHENYLEPHRINE HCl

Nasal decongestant

Relieves:

Sinus headache,
Sinus pressure, Chest
congestion, Nasal
congestion & Mucus



Actual Size

24 CAPLETS

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg Pain reliever/fever reducer
Guaifenesin 200 mg Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Drug Facts (continued)

Uses

temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
■ headache
■ minor aches and pains
■ sinus congestion and pressure
■ temporarily reduces fever

Drug Facts (continued)

■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ rashes
■ skin redness
■ itchy skin
If a skin reaction occurs, stop use and seek medical help right away.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ heart disease
■ thyroid disease
■ diabetes
■ high blood pressure
■ difficulty in urination due to enlargement of the prostate gland
■ cough that occurs with too much phlegm (mucus)
■ liver disease

Drug Facts (continued)

■ see end flap for expiration date and lot number

Inactive ingredients corn starch, croscollon, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if:
■ nervousness, dizziness, or sleeplessness occur
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed
■ adults and children 12 years and over:
■ take 2 caplets every 4 hours
■ swallow whole - do not crush, chew, or dissolve
■ do not take more than 10 caplets in 24 hours
■ children under 12 years: ask a doctor

Other information

■ each caplet contains: sodium 3 mg
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BUSTERS ARE BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Drug Facts (continued)

polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments? 1-800-426-9391

No Print/No Varnish
Lot and Expiration No.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BUSTERS ARE BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844
REV0922052708
Trademark: Tylenol® Sinus Severe.
Corporation, owner of the registered

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-19849



B-0231-527C-08SCHR
REV0922052708



Severe Tussin CF
Non-Drowsy
MAXIMUM STRENGTH
CVS Health

DO NOT USE IF PRINTED NECKBAND
IS BROKEN OR MISSING

CVS Health

Compare to the active ingredients in Robitussin® Severe Multi-Symptom Cough Cold + Flu*

MAXIMUM STRENGTH
NDC 69842-565-26

Non-Drowsy Severe Tussin CF
ACETAMINOPHEN
Pain Reliever/Fever Reducer
DEXTROMETHORPHAN HBr
Cough suppressant
GUAIFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant
Adult Cough & Cold

Relieves:

- Cough, sore throat
- Body aches, fever
- Nasal congestion
- Chest congestion



For Ages 12 & Over
Dosage cup provided
Package Contains One Bottle

4 FL OZ (118 mL)

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen, USP 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr, USP 20 mg.....Cough suppressant
Guaifenesin, USP 400 mg.....Expectorant
Phenylephrine HCl, USP 10 mg.....Nasal decongestant

Uses

- temporarily relieves these symptoms occurring with a cold or flu:
 - cough due to minor throat and bronchial irritation
 - nasal congestion
 - sore throat
 - sinus congestion and pressure
 - minor aches and pains
 - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than 5 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.

Drug Facts (continued)

- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information

- each 20 mL contains: sodium 15 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, benzyl alcohol, edetate disodium, FD&C red #40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, triacetin, xanthan gum

Questions or comments?
1-800-719-9260

*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Severe Multi-Symptom Cough Cold + Flu.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2018 CVS/pharmacy
CVS.com® 1-800-SHOP CVS

V-16430

CVS Quality
Money Back Guarantee

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

#971824



0 50428 31312 1

LOT NO.

EXP.

7C126 17 C1



Drug Facts (continued)	
Uses	<ul style="list-style-type: none"> Temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold: nasal congestion headache runny nose and sneezing sinus congestion and pressure promotes sinus drainage helps clear nasal passages helps decongest sinus openings and passages temporarily reduces fever
Purpose	<p>(Daytime Sinus Congestion & Pain) Pain reliever/fever reducer Acetaminophen 325 mg Chlorpheniramine maleate 2 mg</p> <p>(Nighttime Sinus Congestion & Pain) Pain reliever/fever reducer Acetaminophen 325 mg Chlorpheniramine maleate 2 mg</p>
Warnings	<p>Phenylephrine HCl 5 mg</p> <p>COMPLETE PRODUCT INFORMATION</p>

Drug Facts (continued)	
Warnings	<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.</p> <p>With other drugs containing acetaminophen</p> <p>or more alcoholic drinks every day while using this product</p> <p>Alcohol: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash</p> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p> <p>If you have ever had an allergic reaction to this product or any of its ingredients</p> <p>With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</p> <p>(MAD) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAD drug. If you do not know if your prescription drug contains an MAD, ask a doctor or pharmacist before taking this product.</p> <p>As a doctor before use if you have</p> <ul style="list-style-type: none"> heart disease high blood pressure thyroid disease diabetes breathing problem such as emphysema or chronic bronchitis glaucoma (Nighttime only) <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking sedatives or tranquilizers (Nighttime only) taking the blood thinning drug warfarin taking sedatives or tranquilizers
Directions	<p>Do not take more than directed</p> <ul style="list-style-type: none"> adults and children 12 years and over take 2 caplets every 4 hours do not crush, chew, or dissolve do not take more than 10 caplets in 24 hours children under 12 years: ask a doctor
Drug Facts (continued)	<p>When using this product</p> <ul style="list-style-type: none"> do not exceed recommended dosage use caution when driving a motor vehicle or operating machinery (Nighttime only) do not use if you are drowsy avoid alcoholic beverages (Nighttime only) Stop use and ask a doctor if fever persists or lasts more than 3 days redness or swelling is present new symptoms occur if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Read each section carefully. Do not take DAYTIME and NIGHTTIME products at the same time.

Drug Facts (continued)	
Other information	<p>■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BUSTERS ARE TORN OR BROKEN</p> <p>■ Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)</p> <p>■ See end flap for expiration date and lot number</p>
Inactive ingredients (Daytime only)	<p>corn starch, croscollon, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide</p>
Inactive ingredients (Nighttime only)	<p>corn starch, croscollon, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, powder, silicon dioxide, sodium starch glycolate, stearic acid, sucrose, talc, titanium dioxide</p>
Questions or comments?	<p>1-800-426-9391</p>
Do not take the Daytime and Nighttime caplets at the same time.	<p>50844 REV0721C45546608</p>

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BUSTERS ARE TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2023 CVS Pharmacy
 CVS.com 1-800-SHOP CVS
 V-19849

100% money back guaranteed.
 CVS.com/returnpolicy

No print/No varnish Lot & Exp date

#35238
 050428125156 0

DAYTIME - Non-Drowsy
Sinus Pain & Congestion
 ACETAMINOPHEN
 Pain reliever, Fever reducer
 PHENYLEPHRINE HCl
 Nasal decongestant

Relieves:
 Headache, Sinus pressure & Nasal congestion

12 CAPLETS
 With cool blast flavor

NIGHTTIME
Sinus Pain & Congestion
 ACETAMINOPHEN
 Pain reliever, Fever reducer
 CHLORPHENIRAMINE MALEATE
 Antihistamine
 PHENYLEPHRINE HCl
 Nasal decongestant

Relieves:
 Fever, Headache, Sinus pressure, Nasal congestion & Runny nose

12 CAPLETS
 With cool blast flavor

24 TOTAL CAPLETS



Compare to the active ingredients
in Tylenol® Sinus + Headache*

Daytime

Non-Drowsy

Sinus Pain & Congestion

ACETAMINOPHEN
Pain reliever, Fever reducer

PHENYLEPHRINE HCl
Nasal decongestant

Relieves:
Sinus headache,
Sinus pressure &
Nasal congestion



Actual Size

24 CAPLETS



Drug Facts

Active ingredients (in each capsule)

acetaminophen 325 mg	Pain reliever/fever reducer
phenylephrine HCl 5 mg	Nasal decongestant

Purpose

Temporarily relieves these symptoms associated with hay fever or other respiratory allergies and the common cold:

- runny nose
- itchy throat
- nasal congestion
- sinus congestion and pressure

Drug Facts (continued)

Uses

Temporarily relieves these symptoms associated with hay fever or other respiratory allergies and the common cold:

- minor aches and pains
- headache
- nasal congestion
- sinus congestion and pressure

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Sho use and ask a doctor if:

- you have diabetes or sleeplessness occur
- pain or nasal congestion lasts more than 7 days
- pain or nasal congestion lasts less than 3 days
- redness or swelling of the face
- other symptoms occur

These could be signs of a serious condition.

If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over
- take 2 capsules every 4 hours
- swallow whole – do not crush, chew, or dissolve
- do not take more than 10 capsules in 24 hours
- children under 12 years, ask a doctor

Other information

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

Store at 25°C (77°F); excursions permitted between 15–30°C (59–86°F) store at room temperature.

Use by expiration date and lot number.

#408566

FPO 80%

UPC# 050428296639

X XXXXXXX XXXXXX X

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2022 CVS Pharmacy
CVS.com® 1-800-SHOP CVS
V-19849

100% money back guaranteed.
[CVS.com/returnpolicy](#)

Lot and Expiration No.

Do Not Print/Vo Vanish

NO PRINT/VO VANISH

LOT AND EXPIRATION NO.

TAMPER EVIDENT: DO NOT USE IF UNIT IS TORN, BROKEN OR IF BLISTER PACKAGE IS OPENED ON IF ALUSTER

ANY SIGNS OF TAMPERING

S08044 REV07/21B46608

This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Tylenol® Sinus + Headache.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ rashes
 ■ rash
 ■ skin redness
 ■ a skin reaction occurs, stop use and seek medical help right away.
 Do not use:
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have:

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure



Sinus PE + Allergy

CHLORPHENIRAMINE MALEATE - Antihistamine
PHENYLEPHRINE HYDROCHLORIDE - Nasal decongestant

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2020 CVS/pharmacy
CVS.com 1-800-SHOP CVS
V-19849



No print/No varnish
Lot & Exp date



CVS Quality
Money Back Guarantee

Drug Facts (continued)

Active ingredients (in each tablet)
Chlorpheniramine maleate 4 mg, Antihistamine
Phenylephrine HCl 10 mg, Nasal decongestant

Purpose
Nasal decongestant

Uses
temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies: runny nose, sneezing, itching of the nose or throat, watery eyes, sinus congestion and pressure

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have: high blood pressure, heart disease, thyroid disease, diabetes, difficulty in urination due to enlargement of the prostate gland, a breathing problem such as emphysema or chronic bronchitis.
When using this product, do not exceed recommended dosage. Excitability may occur, especially in children.

Directions
adults and children 12 years and over: take 1 tablet every 4 hours. Do not take more than 6 tablets in 24 hours.
children under 12 years: do not use

Other information
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
see end flap for expiration date and lot number

Inactive ingredients
croscarmellose sodium, lactose anhydrous, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments? 1-800-426-9391

#477524



TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844 REV0319A46208



Sinus PE + Allergy

CHLORPHENIRAMINE MALEATE - Antihistamine
PHENYLEPHRINE HYDROCHLORIDE - Nasal decongestant



Sinus PE + Allergy

CHLORPHENIRAMINE MALEATE - Antihistamine
PHENYLEPHRINE HYDROCHLORIDE - Nasal decongestant

MAXIMUM STRENGTH

Relieves:

- Sinus pressure & Congestion
- Sneezing
- Itchy eyes
- Runny nose

24 TABLETS UNCOATED



Actual Size



B-0231-462-08-H
REV0319A46208



TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® SINUS-MAX® Night Time Congestion & Cough.

#547736



PLD-A378C LB007740

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2020 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-13114



PEEL CORNER FOR DRUG FACTS

Drug Facts**Active ingredients
(in each 20 mL)****Purposes**

Acetaminophen 650 mg..... Pain reliever
Diphenhydramine HCl 25 mg.....
.....Antihistamine/cough suppressant
Phenylephrine HCl 10 mg.....Nasal decongestant

Uses

- temporarily relieves
- nasal congestion ■ headache
- minor aches and pains ■ cough
- sinus congestion and pressure
- runny nose and sneezing
- temporarily promotes nasal and/or sinus drainage
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease ■ heart disease
- high blood pressure ■ diabetes
- thyroid disease ■ glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)**Ask a doctor or pharmacist before use if**

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or

Drug Facts (continued)

contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter ■ keep dosing cup with product
- dose as follows or as directed by a doctor
- adults and children 12 years and older:
20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

- with any other product containing diphenhydramine, even one used on skin (**Nighttime only**)

Ask a doctor before use if you have

- heart disease ■ high blood pressure
- liver disease ■ thyroid disease ■ diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis (**Nighttime only**)
- glaucoma (**Nighttime only**)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**Nighttime only**)

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children (**Nighttime only**)
- marked drowsiness may occur (**Nighttime only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Nighttime only**)

Drug Facts (continued)

- be careful when driving a motor vehicle or operating machinery (**Nighttime only**)
- avoid alcoholic beverages (**Nighttime only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If taking NIGHTTIME and DAYTIME products, carefully read each section to ensure correct dosing. Do not take DAY & NIGHT at the same time.

Drug Facts (continued)

Directions

- do not use more than directed
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients (Daytime only)

corn starch, croscarmellose sodium, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Drug Facts (continued)

Inactive ingredients (Nighttime only)

corn starch, croscarmellose sodium, crospovidone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide

Questions or comments? 1-800-426-9391

50844
ORG051761569409
R-0231-61569409SMHDF

Drug Facts

Active ingredients (in each caplet)

Ingredient	Purpose
Acetaminophen 325 mg	Pain reliever
Guafenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Active ingredients (in each caplet)

Ingredient	Purpose
Acetaminophen 325 mg	Pain reliever
Diphenhydramine HCl 12.5 mg	Antihistamine/cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves:
 - nasal congestion ■ headache ■ minor aches and pains
 - sinus congestion and pressure ■ cough (**Nighttime only**)
 - runny nose and sneezing (**Nighttime only**)
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (**Daytime only**)

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

PEEL HERE FOR MORE DRUG FACTS

CVS Health

Daytime - Sinus Pain
MAXIMUM STRENGTH Sinus Relief

CVS Health

Nighttime - Sinus Pain
MAXIMUM STRENGTH Sinus Relief

#938778
0 50428 26191 0
No print/No varnish area
Lot no/Exp date

CVS Health

Daytime
MAXIMUM STRENGTH Sinus Relief

ACETAMINOPHEN - Pain reliever
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

Relieves:
Sinus pressure & Congestion, Headache
Thins & Loosens mucus

 Actual Size

10 CAPLETS For Ages 12+

CVS Health

Nighttime
MAXIMUM STRENGTH Sinus Relief

ACETAMINOPHEN - Pain reliever
DIPHENHYDRAMINE HCl - Antihistamine/
Cough suppressant
PHENYLEPHRINE HCl - Nasal decongestant

Relieves:
Nasal congestion, Sinus pressure & Pain, Runny nose, Sneezing & Cough

 Actual Size

10 CAPLETS For Ages 12+

Compare to the active ingredients in Maximum Strength Mucinex® SINUS-MAX® Day & Night®

Sinus Pain
NDC 69842-882-09

CVS Health Compare to the active ingredients in Maximum Strength Mucinex® SINUS-MAX® Severe Congestion Relief*
NDC 69842-478-06

ADULT
Severe Congestion Relief
ACETAMINOPHEN -
Pain reliever
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl -
Nasal decongestant
MAXIMUM STRENGTH

Multi-Symptom

- Clears sinus congestion
- Relieves headache
- Thins & loosens mucus

For Ages 12 & Over

6 FL OZ (177 mL)

PLD-A388A LB005809

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® SINUS-MAX® Severe Congestion Relief.

#425051



Distributed by: **CVS Pharmacy, Inc.**
One CVS Drive, Woonsocket, RI 02895
© 2018 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-13114



PLD-A388A LB005809

Drug Facts**Active ingredients
(in each 20 mL)****Purposes**

Acetaminophen 650 mg.....	Pain reliever
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)**Uses**

- temporarily relieves
 - headache
 - nasal congestion
 - minor aches and pains
 - sinus congestion and pressure
 - temporarily promotes nasal and/or sinus drainage
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Drug Facts (continued)**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)**Directions**

- do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday
9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS

<p>Do not use for asthma or children.</p> <p>Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</p> <p>If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>If you ever had an allergic reaction to this product or any of its ingredients, in children 12 years of age.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none">liver diseaseheart diseasehigh blood pressurethyroid diseasediabetescough with excess phlegm (mucus)a breathing problem such as emphysema or chronic bronchitisdifficulty in urination due to enlargement of the prostate glandperforated or drained cough occurs with smoking, asthma, or emphysema <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none">taking sedatives or tranquilizerstaking the blood thinning drug warfarin <p>When using this product do not exceed recommended dosage</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none">fever gets worse or lasts more than 3 daysrunny nose, cough, or nasal congestion gets worse or lasts more than 7 days	<p>Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</p> <p>If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>If you ever had an allergic reaction to this product or any of its ingredients, in children 12 years of age.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none">liver diseaseheart diseasehigh blood pressurethyroid diseasediabetescough with excess phlegm (mucus)difficulty in urination due to enlargement of the prostate glandperforated or drained cough occurs with smoking, asthma, or emphysema <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none">taking the blood thinning drug warfarin <p>When using this product do not exceed recommended dosage</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none">fever gets worse or lasts more than 3 daysrunny nose, cough, or nasal congestion gets worse or lasts more than 7 days
---	---

Distributed by CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2017 CVS/Pharmacy
 CVS.com • 1-800-SHOP CVS
 Made in India
 V-32577
CVS Quality
 Money Back Guarantee
 R-00

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Alike-Salter PILLS® Day & Night Severe Sinus Congestion & Cough Liquid Gels.

CVS Health

Important: Read all product information before using. Keep the carton for important information.
 DO NOT USE IF BLISTER IS CUT OR BROKEN.

Severe Sinus Congestion & Cough Relief
Day

8 SOFTGELS
 Actual Size



Relieves:
 Sinus Congestion
 Headache & Pain,
 Sneezing, Cough,
 Runny Nose &
 Pressure.

Relieves:
 Nasal decongestant
 PHENYLEPHRINE HCl 5 mg each
 Antihistamine
 DOXYLAMINE SUCCINATE 6.25 mg each
 Cough suppressant
 DEXTROMETHORPHAN HBr 10 mg each
 Pain reliever - Fever reducer
 ACETAMINOPHEN 325 mg each

TOTAL 20 SOFTGELS

Severe Sinus Congestion & Cough Relief
Night

12 SOFTGELS
 Actual Size



Relieves:
 Sinus Congestion,
 Headache &
 Body Ache,
 Cough

Relieves:
 Nasal decongestant
 PHENYLEPHRINE HCl 5 mg each
 Cough suppressant
 DEXTROMETHORPHAN HBr 10 mg each
 Pain reliever - Fever reducer
 ACETAMINOPHEN 325 mg each

Severe Sinus Congestion & Cough Relief Day

Drug Facts

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Phenylephrine hydrochloride 5 mg.....	Nasal decongestant

Uses

- temporarily relieves these symptoms due to cold and flu:
 - sneezing
 - itching of the nose, throat or watery eyes due to hay fever
 - cough
 - nasal congestion
 - sinus congestion and pressure
 - sore throat
 - headache
 - minor aches and pains
- helps clear nasal passages and shrinks swollen membranes
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 10 softgels in 24 hours, which is the maximum daily amount for this product.
- With other drugs containing acetaminophen.
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- hives
- skin redness
- blister
- rash
- hives

Severe Sinus Congestion & Cough Relief Night

Drug Facts

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine
Phenylephrine hydrochloride 5 mg.....	Nasal decongestant

Uses

- temporarily relieves these symptoms due to cold and flu:
 - sneezing
 - itching of the nose, throat or watery eyes due to hay fever
 - cough
 - nasal congestion
 - sinus congestion and pressure
 - sore throat
 - headache
 - minor aches and pains
 - runny nose
- helps clear nasal passages and shrinks swollen membranes
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 10 softgels in 24 hours, which is the maximum daily amount for this product.
- With other drugs containing acetaminophen.
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- hives
- skin redness
- blister
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

Severe Sinus Congestion & Cough Relief Day

Drug Facts (continued)

- facial swelling
- asthma (wheezing)
- shock

If a skin or allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting consult a doctor promptly.

Severe Sinus Congestion & Cough Relief Night


Drug Facts (continued)

If a skin or allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting consult a doctor promptly.

Exp. Lot

COATING PEEF AREA



#234944

TAMPER EVIDENT: DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredients (in each 10 mL)

Brompheniramine Maleate, USP 2 mg.....Antihistamine
Phenylephrine HCl, USP 5 mg.....Nasal Decongestant

Purposes

- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily restores freer breathing through the nose.

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem such as emphysema, asthma, or chronic bronchitis.

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- ml = milliliter

age	dose
adults and children 12 years and over	20 ml every 4 hours
children 6 to under 12 years	10 ml every 4 hours
children under 6 years	do not use.

Other information

- each 10 ml contains: sodium 6 mg
- store at room temperature.

Inactive ingredients citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sodium citrate, sorbitol liquid, sucralose.

Questions or comments? 1-866-467-2748

CVS Health® Children's Cold & Allergy grape flavor offers relief from nasal congestion, stuffy nose, sneezing, runny nose, itchy, watery eyes, and itchy nose and throat to get through the day.

CVS Health. Grape Flavor

Children's Cold & Allergy Liquid

Compare to the active ingredients in Children's Dimetapp® Cold & Allergy*.

CVS Health. Grape Flavor

Children's Cold & Allergy

BROMPHENIRAMINE MALEATE, USP
Antihistamine
PHENYLEPHRINE HCl, USP
Nasal decongestant

Liquid

Relieves:
Nasal congestion; Stuffy nose; Runny nose; Sneezing; Itchy, watery eyes; Itchy nose & throat

Alcohol free

Actual Bottle Size on Side Panel



Grape Flavor
For Ages 6 Years & Over

4 FL OZ (118 mL)

#477078



69304CVSFC 0 50428 25668 8

Lot:

Exp:

IMPORTANT:
Keep this carton for future reference on full labeling.

Package Contains
One Bottle

Actual Size

*CVS Health® Children's Cold & Allergy is not manufactured or distributed by Pfizer, distributor of Children's Dimetapp® Cold & Allergy.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2020 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-12431

CVS Quality
Money Back Guarantee

10

DAYTIME SEVERE COLD AND FLU	NIGHTTIME SEVERE COLD AND FLU
Drug Facts (continued) How throat lozenges work: Throat lozenges provide relief for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Warnings: Acetaminophen may cause severe liver problems. Symptoms may include: <input type="checkbox"/> skin redness <input type="checkbox"/> itchy skin <input type="checkbox"/> rash <input type="checkbox"/> a skin reaction occurs, stop use and seek medical help right away. Do not use: <input type="checkbox"/> With any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <input type="checkbox"/> If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, Parkinson's, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have: <input type="checkbox"/> Heart disease <input type="checkbox"/> heart disease <input type="checkbox"/> high blood pressure <input type="checkbox"/> Peptic disease <input type="checkbox"/> diabetes <input type="checkbox"/> asthma <input type="checkbox"/> trouble swallowing due to enlarged prostate gland <input type="checkbox"/> cough that occurs with too much phlegm (mucus) <input type="checkbox"/> persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema. Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. When using this product, do not use more than directed. May use and ask a doctor if: <input type="checkbox"/> you get nervous, dizzy or sleepless <input type="checkbox"/> you have constipation, or cough gets worse or lasts more than 7 days	Drug Facts (continued) Very alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <input type="checkbox"/> skin redness <input type="checkbox"/> itchy skin <input type="checkbox"/> rash <input type="checkbox"/> a skin reaction occurs, stop use and seek medical help right away. Do not use: <input type="checkbox"/> With any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <input type="checkbox"/> If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, Parkinson's, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Do not take a sleep aid: Ask a doctor before use if you have: <input type="checkbox"/> Heart disease <input type="checkbox"/> heart disease <input type="checkbox"/> high blood pressure <input type="checkbox"/> Peptic disease <input type="checkbox"/> diabetes <input type="checkbox"/> asthma <input type="checkbox"/> cough that occurs with too much phlegm (mucus) <input type="checkbox"/> a breathing problem or chronic cough that lasts or so occurs with smoking, asthma, chronic bronchitis, or emphysema <input type="checkbox"/> trouble swallowing due to enlarged prostate gland Ask a doctor or pharmacist before use if you are: <input type="checkbox"/> taking sedatives or tranquilizers <input type="checkbox"/> taking the blood thinning drug warfarin When using this product: <input type="checkbox"/> Do not use more than directed <input type="checkbox"/> Avoidance may occur, especially in children <input type="checkbox"/> avoid alcohol drinks



READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.
KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

Drug Facts

Active Ingredients (in each packet)

Acetaminophen 500 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 20 mg	Cough suppressant
Phenylephrine hydrochloride 10 mg	Nasal Decongestant

Uses

- Temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
 - temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.

Do not use if you are in a child under 4 years of age.

Do not use if you are allergic to acetaminophen or with any other drug containing acetaminophen (prescription and nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are:

- taking the blood thinning drug warfarin

When using this product:

- Do not exceed recommended dosage

Stop use and ask a doctor if:

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

*This product is not manufactured or distributed by Novartis Consumer Health, Inc., distributor of Theraflu® Multi-Symptom Severe Cold.

Drug Facts (continued)

Directions

- Do not use more than directed
- Take every 4 hours, not to exceed 6 packets in 24 hours or as directed by a doctor

Age	Dose
children under 4 years of age	do not use
children 4 to under 12 years of age	do not use unless directed by a doctor
adults and children 12 years of age and over	one packet

■ dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.

■ If using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 6 mg
- store at controlled room temperature 20°-25°C (68°-77°F). Protect from excessive heat and moisture.

Inactive ingredients anhydrous citric acid, caramel, flavors, potassium citrate, silica, sucralose, sucrose, xanthan gum

Questions or comments?
1-866-467-2748

TAMPER EVIDENT INNER UNIT:
DO NOT USE IF SEALED PACKET IS TORN OR BROKEN.

Dye, Aspartame, & Sodium free

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2015 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-12431

CVS Quality
Money Back Guarantee

PLEASE RECYCLE

CVS Health.

Multi-Symptom Severe Cold

ACETAMINOPHEN
Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

6 PACKETS

Compare to the active ingredients in Theraflu® Multi-Symptom Severe Cold®

Severe

NDC 59779-877-06

Multi-Symptom Severe Cold

ACETAMINOPHEN
Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:

- Nasal congestion
- Sore throat pain
- Cough
- Headache
- Body ache
- Fever

Green Tea & Honey Lemon Flavors

6 PACKETS

CVS Health.

Multi-Symptom Severe Cold

CVS Health™ Multi-Symptom Severe Cold provides complete relief of your severe cold symptoms. This soothing hot liquid formula is dye free, aspartame free and sodium free and tastes great.

For nighttime relief, try CVS Health™ Nighttime Severe Cough & Cold.



#633103 54706CVSFC

0 50428 52409 1


Lot:

Exp:



FRONT**BACK-1****3 COLOR
MAX - ART**FPO 80%
UPC# 050428300985**BACK-2****1 COLOR
MAX - ART****BACK-3****1 COLOR
MAX - ART**

POWERFUL
NIGHT TIME
RELIEF





Drug Facts (continued)

Severe Sinus Congestion & Cough Day

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough with excessive phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Drug Facts (continued)

Severe Sinus Congestion & Cough Night

Do not use to sedate children.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes ■ glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- may cause marked drowsiness ■ avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children



COLD & FLU FORMULA • DAY NON-DROWSY**Drug Facts** (continued)

Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

■ in children under 12 years of age

Ask a doctor before use if you have

■ liver disease ■ heart disease ■ high blood pressure

■ thyroid disease ■ diabetes

■ cough with excessive phlegm (mucus)

■ difficulty in urination due to enlargement of the prostate gland

■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage

COLD & FLU FORMULA • NIGHT**Drug Facts** (continued)**Do not use**

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

■ in children under 12 years of age

Ask a doctor before use if you have

■ liver disease ■ heart disease ■ high blood pressure

■ thyroid disease ■ diabetes ■ glaucoma

■ cough with excessive phlegm (mucus)

■ a breathing problem such as emphysema or chronic bronchitis

■ difficulty in urination due to enlargement of the prostate gland

■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

■ taking the blood thinning drug warfarin

■ taking sedatives or tranquilizers

When using this product

■ **do not exceed recommended dosage**

■ may cause marked drowsiness ■ avoid alcoholic drinks

■ alcohol, sedatives, and tranquilizers may increase drowsiness

■ be careful when driving a motor vehicle or operating machinery

■ excitability may occur, especially in children

Drug Facts

Active ingredients (in each 15 mL) Purposes

Acetaminophen 325 mg.....Pain reliever/fever reducer

Dextromethorphan HBr 10 mg.....Cough suppressant

Phenylephrine HCl 5 mg.....Nasal decongestant

Uses ■ temporarily relieves common cold and flu symptoms

■ minor aches and pains ■ headache ■ sore throat

■ nasal congestion ■ fever

■ cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if: ■ adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount ■ child takes more than 4 doses (15 mL each) in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

← PEEL HERE: **Drug Facts** (continued under label)

CVS Health.

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu*

NDC 51316-486-08

Non-Drowsy Daytime

ACETAMINOPHEN
Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr
Cough suppressant

PHENYLEPHRINE HCl
Nasal decongestant

Relieves:
Headache, Fever, Sore throat, Minor aches & pains, Nasal congestion, Sinus pressure, Cough

COLD & FLU

Antihistamine free
Ages 6+

8 FL OZ (237 mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02896
© 2023 CVS/pharmacy
CVS.com®
1-800-SHOP CVS
V-16425

100% money back guaranteed.
CVS.com/returnpolicy

#890115

0 50428 29666 0

PLD-A3800 LB009123

Drug Facts (continued)

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease

■ heart disease ■ high blood pressure ■ thyroid disease

■ diabetes ■ a sodium-restricted diet ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 days (adult) ■ fever gets worse, or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended

Drug Facts (continued)

dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in any 24-hour period ■ measure only with dosing cup provided. Do not use any other dosing device.

■ keep dosing cup with product ■ mL = milliliter

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information


■ each 15 mL contains: sodium 12 mg

■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

*



KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg.....	Pain reliever/Fever reducer
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

- temporarily relieves symptoms due to the common cold
 - minor aches and pains
 - headache
 - fever
 - nasal congestion
 - sinus congestion & pressure
- temporarily relieves symptoms due to hay fever or other upper respiratory allergies
 - minor aches and pains
 - headache
 - nasal congestion
 - sinus congestion & pressure

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hours, which is the maximum daily amount for this product.

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- you get nervous, dizzy or sleepless

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at room temperature 15°-30°C (59°-86°F)

Drug Facts (continued)

Inactive ingredients

FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution and white ink

Questions or comments? Call toll free: 1-855-215-8180

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2017 CVS/pharmacy
CVS.com® 1-800-SHOP CVS

Made in China
V-31177



CVS27-00

*This product is not manufactured or distributed by The Procter & Gamble Manufacturing Company, owner of the registered trademarks Vicks® and Sinex™ Daytime Congestion Pressure & Pain.

2697

Drug Facts

Active Ingredients

Aspirin 325 mg (NSAID)* Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg Antihistamine
Phenylephrine bitartrate 7.5 mg Nasal decongestant

Uses

temporarily relieves these symptoms of the common cold:
■ fever ■ sneezing ■ nasal congestion ■ sore throat
■ headache ■ minor aches and pains ■ runny nose

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using the product ■ take more or for a longer time than directed

Do not use

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you are allergic to aspirin ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or medical conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if ■ you have a history of stomach problems, such as heartburn ■ you are taking blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have ■ you are taking a diuretic ■ you are taking sedatives or tranquilizers ■ you have ■ you have a gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

Ask a doctor or pharmacist before use if you are ■ presently taking a prescription drug ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by doctor.

Do not use more than directed ■ you may get drowsy ■ avoid alcoholic drinks ■ excitability may occur, especially in children ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ an allergic reaction occurs.

Drug Facts (continued)

Sore throat warning: If sore throat is severe, headache, rash, nausea, or vomiting, accompanied or followed by fever, consult a doctor promptly.

CVS
Health

Orange Flavor

Effervescent Cold Relief



Actual Size

CVS
Health

Compare to the active ingredients in Alka-Seltzer Plus® Cold Orange Zest†

Orange Flavor

NDC 69842-875-20

Effervescent Cold Relief

ASPIRIN (NSAID) - Pain reliever; Fever reducer
CHLORPHENIRAMINE MALEATE - Antihistamine
PHENYLEPHRINE BITARTRATE
Nasal decongestant

Fast relief of:

Nasal congestion; Runny nose; Sneezing; Headache; Body ache; Sore throat; Sinus pain & pressure



Orange Flavor

20 TABLETS

Actual Size Tablet on Top Panel



Drug Facts (continued)

Seek medical help right away.
■ pain or nasal congestion gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days ■ new symptoms occur
■ redness or swelling is present
■ ringing in the ears or a loss of hearing occurs ■ nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not use more than directed (see overdose warning)
■ adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4-6 hours
■ do not take more than 8 tablets in 24 hours
■ children under 12 years: ask a doctor

Drug Facts (continued)

Other information

each tablet contains:
sodium 464 mg
■ phenylketonurics: contains phenylalanine 9 mg per tablet
■ store at room temperature (59°-86°F)

Inactive ingredients

acesulfame potassium, aspartame, citric acid, docusate sodium, FD&C red #40, FD&C yellow #6, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2017 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-17432



†This product is not manufactured, distributed or licensed by the maker of Alka-Seltzer Plus® Cold Orange Zest®.

#182122



DO NOT USE IF PRINTED
PACKETS ARE TORN OR PUNCTURED

PN: 87560 CH C1
REV: 02/17

CVS Health.
MAXIMUM STRENGTH
Cold & Flu

*This product is not manufactured or distributed by Reckitt Benckiser LLC, owner of the registered trademark Maximum Strength Mucinex® Fast-Max® Cold & Flu, 6281 0421

SAFETY SEALED: DO NOT USE IF INDIVIDUAL BUSTLE UNIT IS OPEN OR TORN.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

Questions? Call 1-877-290-4008

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2021 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
Made in India V-37828

Drug Facts

Active ingredients (in each softgel)

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- Temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - headache
 - cough
 - minor aches & pains
 - sore throat
- Temporarily reduces fever
- Helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 softgels in 24 hrs, which is the maximum daily amount;
- with other drugs containing acetaminophen; 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have: liver disease, heart disease, diabetes, high blood pressure, thyroid disease, trouble urinating due to enlarged prostate gland, persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, or cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if: nervousness, dizziness, or sleeplessness occur; pain, nasal congestion, or cough gets worse or lasts more than 7 days; liver gets worse or lasts more than 3 days; redness or swelling is present; new symptoms occur; cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

CVS Health.

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® Cold & Flu®
NDC 69842-789-16

MAXIMUM STRENGTH

Fast Dissolving SoftGels

Cold & Flu

ACETAMINOPHEN - Pain reliever / Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

MULTI-SYMPOM

- Controls cough
- Thins & loosens mucus
- Relieves nasal & chest congestion
- Relieves aches, fever & sore throat

FOR AGES 12+
Actual Size

16 SOFTGELS† (†Liquid-filled capsules)

Print/Varnish Omit Area

CVS Health.
MAXIMUM STRENGTH
Cold & Flu

*This product is not manufactured or distributed by Reckitt Benckiser LLC, owner of the registered trademark Maximum Strength Mucinex® Fast-Max® Cold & Flu, 6281 0421

SAFETY SEALED: DO NOT USE IF INDIVIDUAL BUSTLE UNIT IS OPEN OR TORN.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

Questions? Call 1-877-290-4008

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2021 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
Made in India V-37828

Drug Facts

Active ingredients (in each softgel)

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- Temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - headache
 - cough
 - minor aches & pains
 - sore throat
- Temporarily reduces fever
- Helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 softgels in 24 hrs, which is the maximum daily amount;
- with other drugs containing acetaminophen; 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have: liver disease, heart disease, diabetes, high blood pressure, thyroid disease, trouble urinating due to enlarged prostate gland, persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, or cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if: nervousness, dizziness, or sleeplessness occur; pain, nasal congestion, or cough gets worse or lasts more than 7 days; liver gets worse or lasts more than 3 days; redness or swelling is present; new symptoms occur; cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

CVS Health.

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® Cold & Flu®
NDC 69842-789-16

MAXIMUM STRENGTH

Fast Dissolving SoftGels

Cold & Flu

ACETAMINOPHEN - Pain reliever / Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

MULTI-SYMPOM

- Controls cough
- Thins & loosens mucus
- Relieves nasal & chest congestion
- Relieves aches, fever & sore throat

FOR AGES 12+
Actual Size

16 SOFTGELS† (†Liquid-filled capsules)

Print/Varnish Omit Area

CVS Health.
MAXIMUM STRENGTH
Cold & Flu

*This product is not manufactured or distributed by Reckitt Benckiser LLC, owner of the registered trademark Maximum Strength Mucinex® Fast-Max® Cold & Flu, 6281 0421

SAFETY SEALED: DO NOT USE IF INDIVIDUAL BUSTLE UNIT IS OPEN OR TORN.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

Questions? Call 1-877-290-4008

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2021 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
Made in India V-37828

Drug Facts

Active ingredients (in each softgel)

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- Temporarily relieves these common cold and flu symptoms:
 - nasal congestion
 - headache
 - cough
 - minor aches & pains
 - sore throat
- Temporarily reduces fever
- Helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 softgels in 24 hrs, which is the maximum daily amount;
- with other drugs containing acetaminophen; 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have: liver disease, heart disease, diabetes, high blood pressure, thyroid disease, trouble urinating due to enlarged prostate gland, persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, or cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if: nervousness, dizziness, or sleeplessness occur; pain, nasal congestion, or cough gets worse or lasts more than 7 days; liver gets worse or lasts more than 3 days; redness or swelling is present; new symptoms occur; cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

CVS Health.

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® Cold & Flu®
NDC 69842-789-16

MAXIMUM STRENGTH

Fast Dissolving SoftGels

Cold & Flu

ACETAMINOPHEN - Pain reliever / Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

MULTI-SYMPOM

- Controls cough
- Thins & loosens mucus
- Relieves nasal & chest congestion
- Relieves aches, fever & sore throat

FOR AGES 12+
Actual Size

16 SOFTGELS† (†Liquid-filled capsules)

Print/Varnish Omit Area

CVS Health

MAXIMUM STRENGTH

Cold, Flu & Sore Throat

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieve these common cold and flu symptoms: nasal congestion, sore throat, sinus congestion and pressure, minor aches & pains, temporarily reduces fever, helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 softgels in 24 hrs, which is the maximum daily amount.
- with other drugs containing acetaminophen.
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have: liver disease, heart disease, diabetes, high blood pressure, thyroid disease, trouble urinating due to enlarged prostate gland, persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, cough that occurs with too much phlegm (mucus).

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if: nervousness, dizziness, or sleeplessness occur; pain, nasal congestion, or cough gets worse or lasts more than 7 days; fever gets worse or lasts more than 3 days; redness or swelling is present; new symptoms occur; cough comes back or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Questions? Call 1-877-290-4008

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woodstock, NJ 02899
© 2021 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
Made in India V-37828

Lot #: _____

Exp. Date: _____

Print/Varnish Omit Area

PARENTS:
Learn about how medicine works
www.StopMedicineNow.org

CVS Health

MAXIMUM STRENGTH

Cold, Flu & Sore Throat

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® Cold, Flu & Sore Throat*

NDC 69842-616-16

MAXIMUM STRENGTH

Cold, Flu & Sore Throat

Fast Dissolving SoftGels

ACETAMINOPHEN - 325 mg Pain reliever / Fever reducer
DEXTROMETHORPHAN HBr - 10 mg Cough suppressant
GUAIFENESIN - 200 mg Expectorant
PHENYLEPHRINE HCl - 5 mg Nasal decongestant

- Controls cough
- Relieves nasal & chest congestion, headache & fever
- Thins & loosens mucus

16 SOFTGELS† (†Liquid-filled capsules)

Print/Varnish Omit Area

***This product is not manufactured or distributed by Reckitt Benckiser LLC, owner of the registered trademark Maximum Strength Mucinex® Fast-Max® Day Cold & Flu and Night Cold & Flu. ©2021 CVS Health.**

SAFETY SEALED: DO NOT USE IF ORIGINAL BLISTER UNIT IS OPEN OR TORN.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2021 CVS/pharmacy
CVS.com* 1-800-SHOP CVS
Made in India V-37828

Varnish Omit Area

DAY & NIGHT COMBO PACK

CVS Health.

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® Day Cold & Flu and Night Cold & Flu
NDC 69842-451-24

MAXIMUM STRENGTH

Daytime Cold & Flu

ACETAMINOPHEN
325 mg Pain reliever / Fever reducer

DEXTROMETHORPHAN HBr
10 mg Cough suppressant

GUAIFENESIN
200 mg Expectorant

PHENYLEPHRINE HCl
5 mg Nasal decongestant

MULTI-SYMPTOM

- Controls cough
- Thins & loosens mucus
- Relieves nasal & chest congestion
- Relieves aches, fever & sore throat

FOR AGES 12+
Actual Size

16 SOFTGELS†
24 TOTAL (†Liquid-filled capsules)

MAXIMUM STRENGTH

Nighttime Cold & Flu

ACETAMINOPHEN
325 mg Pain reliever / Fever reducer

DEXTROMETHORPHAN HBr
10 mg Cough suppressant

DOXYLAMINE SUCCINATE
6.25 mg Antihistamine

PHENYLEPHRINE HCl
5 mg Nasal decongestant

MULTI-SYMPTOM

- Controls cough
- Relieves runny nose & sneezing
- Relieves nasal congestion
- Relieves aches, fever & sore throat

FOR AGES 12+
Actual Size

8 SOFTGELS†

Print/Varnish Omit Area

Drug Facts (continued)

Other Ingredients (continued)
FD-302, FD-303, FD-304, FD-305, FD-306, FD-307, FD-308, FD-309, FD-310, FD-311, FD-312, FD-313, FD-314, FD-315, FD-316, FD-317, FD-318, FD-319, FD-320, FD-321, FD-322, FD-323, FD-324, FD-325, FD-326, FD-327, FD-328, FD-329, FD-330, FD-331, FD-332, FD-333, FD-334, FD-335, FD-336, FD-337, FD-338, FD-339, FD-340, FD-341, FD-342, FD-343, FD-344, FD-345, FD-346, FD-347, FD-348, FD-349, FD-350, FD-351, FD-352, FD-353, FD-354, FD-355, FD-356, FD-357, FD-358, FD-359, FD-360, FD-361, FD-362, FD-363, FD-364, FD-365, FD-366, FD-367, FD-368, FD-369, FD-370, FD-371, FD-372, FD-373, FD-374, FD-375, FD-376, FD-377, FD-378, FD-379, FD-380, FD-381, FD-382, FD-383, FD-384, FD-385, FD-386, FD-387, FD-388, FD-389, FD-390, FD-391, FD-392, FD-393, FD-394, FD-395, FD-396, FD-397, FD-398, FD-399, FD-400, FD-401, FD-402, FD-403, FD-404, FD-405, FD-406, FD-407, FD-408, FD-409, FD-410, FD-411, FD-412, FD-413, FD-414, FD-415, FD-416, FD-417, FD-418, FD-419, FD-420, FD-421, FD-422, FD-423, FD-424, FD-425, FD-426, FD-427, FD-428, FD-429, FD-430, FD-431, FD-432, FD-433, FD-434, FD-435, FD-436, FD-437, FD-438, FD-439, FD-440, FD-441, FD-442, FD-443, FD-444, FD-445, FD-446, FD-447, FD-448, FD-449, FD-450, FD-451, FD-452, FD-453, FD-454, FD-455, FD-456, FD-457, FD-458, FD-459, FD-460, FD-461, FD-462, FD-463, FD-464, FD-465, FD-466, FD-467, FD-468, FD-469, FD-470, FD-471, FD-472, FD-473, FD-474, FD-475, FD-476, FD-477, FD-478, FD-479, FD-480, FD-481, FD-482, FD-483, FD-484, FD-485, FD-486, FD-487, FD-488, FD-489, FD-490, FD-491, FD-492, FD-493, FD-494, FD-495, FD-496, FD-497, FD-498, FD-499, FD-500, FD-501, FD-502, FD-503, FD-504, FD-505, FD-506, FD-507, FD-508, FD-509, FD-510, FD-511, FD-512, FD-513, FD-514, FD-515, FD-516, FD-517, FD-518, FD-519, FD-520, FD-521, FD-522, FD-523, FD-524, FD-525, FD-526, FD-527, FD-528, FD-529, FD-530, FD-531, FD-532, FD-533, FD-534, FD-535, FD-536, FD-537, FD-538, FD-539, FD-540, FD-541, FD-542, FD-543, FD-544, FD-545, FD-546, FD-547, FD-548, FD-549, FD-550, FD-551, FD-552, FD-553, FD-554, FD-555, FD-556, FD-557, FD-558, FD-559, FD-560, FD-561, FD-562, FD-563, FD-564, FD-565, FD-566, FD-567, FD-568, FD-569, FD-570, FD-571, FD-572, FD-573, FD-574, FD-575, FD-576, FD-577, FD-578, FD-579, FD-580, FD-581, FD-582, FD-583, FD-584, FD-585, FD-586, FD-587, FD-588, FD-589, FD-590, FD-591, FD-592, FD-593, FD-594, FD-595, FD-596, FD-597, FD-598, FD-599, FD-600, FD-601, FD-602, FD-603, FD-604, FD-605, FD-606, FD-607, FD-608, FD-609, FD-610, FD-611, FD-612, FD-613, FD-614, FD-615, FD-616, FD-617, FD-618, FD-619, FD-620, FD-621, FD-622, FD-623, FD-624, FD-625, FD-626, FD-627, FD-628, FD-629, FD-630, FD-631, FD-632, FD-633, FD-634, FD-635, FD-636, FD-637, FD-638, FD-639, FD-640, FD-641, FD-642, FD-643, FD-644, FD-645, FD-646, FD-647, FD-648, FD-649, FD-650, FD-651, FD-652, FD-653, FD-654, FD-655, FD-656, FD-657, FD-658, FD-659, FD-660, FD-661, FD-662, FD-663, FD-664, FD-665, FD-666, FD-667, FD-668, FD-669, FD-670, FD-671, FD-672, FD-673, FD-674, FD-675, FD-676, FD-677, FD-678, FD-679, FD-680, FD-681, FD-682, FD-683, FD-684, FD-685, FD-686, FD-687, FD-688, FD-689, FD-690, FD-691, FD-692, FD-693, FD-694, FD-695, FD-696, FD-697, FD-698, FD-699, FD-700, FD-701, FD-702, FD-703, FD-704, FD-705, FD-706, FD-707, FD-708, FD-709, FD-710, FD-711, FD-712, FD-713, FD-714, FD-715, FD-716, FD-717, FD-718, FD-719, FD-720, FD-721, FD-722, FD-723, FD-724, FD-725, FD-726, FD-727, FD-728, FD-729, FD-730, FD-731, FD-732, FD-733, FD-734, FD-735, FD-736, FD-737, FD-738, FD-739, FD-740, FD-741, FD-742, FD-743, FD-744, FD-745, FD-746, FD-747, FD-748, FD-749, FD-750, FD-751, FD-752, FD-753, FD-754, FD-755, FD-756, FD-757, FD-758, FD-759, FD-760, FD-761, FD-762, FD-763, FD-764, FD-765, FD-766, FD-767, FD-768, FD-769, FD-770, FD-771, FD-772, FD-773, FD-774, FD-775, FD-776, FD-777, FD-778, FD-779, FD-780, FD-781, FD-782, FD-783, FD-784, FD-785, FD-786, FD-787, FD-788, FD-789, FD-790, FD-791, FD-792, FD-793, FD-794, FD-795, FD-796, FD-797, FD-798, FD-799, FD-800, FD-801, FD-802, FD-803, FD-804, FD-805, FD-806, FD-807, FD-808, FD-809, FD-810, FD-811, FD-812, FD-813, FD-814, FD-815, FD-816, FD-817, FD-818, FD-819, FD-820, FD-821, FD-822, FD-823, FD-824, FD-825, FD-826, FD-827, FD-828, FD-829, FD-830, FD-831, FD-832, FD-833, FD-834, FD-835, FD-836, FD-837, FD-838, FD-839, FD-840, FD-841, FD-842, FD-843, FD-844, FD-845, FD-846, FD-847, FD-848, FD-849, FD-850, FD-851, FD-852, FD-853, FD-854, FD-855, FD-856, FD-857, FD-858, FD-859, FD-860, FD-861, FD-862, FD-863, FD-864, FD-865, FD-866, FD-867, FD-868, FD-869, FD-870, FD-871, FD-872, FD-873, FD-874, FD-875, FD-876, FD-877, FD-878, FD-879, FD-880, FD-881, FD-882, FD-883, FD-884, FD-885, FD-886, FD-887, FD-888, FD-889, FD-890, FD-891, FD-892, FD-893, FD-894, FD-895, FD-896, FD-897, FD-898, FD-899, FD-900, FD-901, FD-902, FD-903, FD-904, FD-905, FD-906, FD-907, FD-908, FD-909, FD-910, FD-911, FD-912, FD-913, FD-914, FD-915, FD-916, FD-917, FD-918, FD-919, FD-920, FD-921, FD-922, FD-923, FD-924, FD-925, FD-926, FD-927, FD-928, FD-929, FD-930, FD-931, FD-932, FD-933, FD-934, FD-935, FD-936, FD-937, FD-938, FD-939, FD-940, FD-941, FD-942, FD-943, FD-944, FD-945, FD-946, FD-947, FD-948, FD-949, FD-950, FD-951, FD-952, FD-953, FD-954, FD-955, FD-956, FD-957, FD-958, FD-959, FD-960, FD-961, FD-962, FD-963, FD-964, FD-965, FD-966, FD-967, FD-968, FD-969, FD-970, FD-971, FD-972, FD-973, FD-974, FD-975, FD-976, FD-977, FD-978, FD-979, FD-980, FD-981, FD-982, FD-983, FD-984, FD-985, FD-986, FD-987, FD-988, FD-989, FD-990, FD-991, FD-992, FD-993, FD-994, FD-995, FD-996, FD-997, FD-998, FD-999, FD-1000, FD-1001, FD-1002, FD-1003, FD-1004, FD-1005, FD-1006, FD-1007, FD-1008, FD-1009, FD-1010, FD-1011, FD-1012, FD-1013, FD-1014, FD-1015, FD-1016, FD-1017, FD-1018, FD-1019, FD-1020, FD-1021, FD-1022, FD-1023, FD-1024, FD-1025, FD-1026, FD-1027, FD-1028, FD-1029, FD-1030, FD-1031, FD-1032, FD-1033, FD-1034, FD-1035, FD-1036, FD-1037, FD-1038, FD-1039, FD-1040, FD-1041, FD-1042, FD-1043, FD-1044, FD-1045, FD-1046, FD-1047, FD-1048, FD-1049, FD-1050, FD-1051, FD-1052, FD-1053, FD-1054, FD-1055, FD-1056, FD-1057, FD-1058, FD-1059, FD-1060, FD-1061, FD-1062, FD-1063, FD-1064, FD-1065, FD-1066, FD-1067, FD-1068, FD-1069, FD-1070, FD-1071, FD-1072, FD-1073, FD-1074, FD-1075, FD-1076, FD-1077, FD-1078, FD-1079, FD-1080, FD-1081, FD-1082, FD-1083, FD-1084, FD-1085, FD-1086, FD-1087, FD-1088, FD-1089, FD-1090, FD-1091, FD-1092, FD-1093, FD-1094, FD-1095, FD-1096, FD-1097, FD-1098, FD-1099, FD-1100, FD-1101, FD-1102, FD-1103, FD-1104, FD-1105, FD-1106, FD-1107, FD-1108, FD-1109, FD-1110, FD-1111, FD-1112, FD-1113, FD-1114, FD-1115, FD-1116, FD-1117, FD-1118, FD-1119, FD-1120, FD-1121, FD-1122, FD-1123, FD-1124, FD-1125, FD-1126, FD-1127, FD-1128, FD-1129, FD-1130, FD-1131, FD-1132, FD-1133, FD-1134, FD-1135, FD-1136, FD-1137, FD-1138, FD-1139, FD-1140, FD-1141, FD-1142, FD-1143, FD-1144, FD-1145, FD-1146, FD-1147, FD-1148, FD-1149, FD-1150, FD-1151, FD-1152, FD-1153, FD-1154, FD-1155, FD-1156, FD-1157, FD-1158, FD-1159, FD-1160, FD-1161, FD-1162, FD-1163, FD-1164, FD-1165, FD-1166, FD-1167, FD-1168, FD-1169, FD-1170, FD-1171, FD-1172, FD-1173, FD-1174, FD-1175, FD-1176, FD-1177, FD-1178, FD-1179, FD-1180, FD-1181, FD-1182, FD-1183, FD-1184, FD-1185, FD-1186, FD-1187, FD-1188, FD-1189, FD-1190, FD-1191, FD-1192, FD-1193, FD-1194, FD-1195, FD-1196, FD-1197, FD-1198, FD-1199, FD-1200, FD-1201, FD-1202, FD-1203, FD-1204, FD-1205, FD-1206, FD-1207, FD-1208, FD-1209, FD-1210, FD-1211, FD-1212, FD-1213, FD-1214, FD-1215, FD-1216, FD-1217, FD-1218, FD-1219, FD-1220, FD-1221, FD-1222, FD-1223, FD-1224, FD-1225, FD-1226, FD-1227, FD-1228, FD-1229, FD-1230, FD-1231, FD-1232, FD-1233, FD-1234, FD-1235, FD-1236, FD-1237, FD-1238, FD-1239, FD-1240, FD-1241, FD-1242, FD-1243, FD-1244, FD-1245, FD-1246, FD-1247, FD-1248, FD-1249, FD-1250, FD-1251, FD-1252, FD-1253, FD-1254, FD-1255, FD-1256, FD-1257, FD-1258, FD-1259, FD-1260, FD-1261, FD-1262, FD-1263, FD-1264, FD-1265, FD-1266, FD-1267, FD-1268, FD-1269, FD-1270, FD-1271, FD-1272, FD-1273, FD-1274, FD-1275, FD-1276, FD-1277, FD-1278, FD-1279, FD-1280, FD-1281, FD-1282, FD-1283, FD-1284, FD-1285, FD-1286, FD-1287, FD-1288, FD-1289, FD-1290, FD-1291, FD-1292, FD-1293, FD-1294, FD-1295, FD-1296, FD-1297, FD-1298, FD-1299, FD-1300, FD-1301, FD-1302, FD-1303, FD-1304, FD-1305, FD-1306, FD-1307, FD-1308, FD-1309, FD-1310, FD-1311, FD-1312, FD-1313, FD-1314, FD-1315, FD-1316, FD-1317, FD-1318, FD-1319, FD-1320, FD-1321, FD-1322, FD-1323, FD-1324, FD-1325, FD-1326, FD-1327, FD-1328, FD-1329, FD-1330, FD-1331, FD-1332, FD-1333, FD-1334, FD-1335, FD-1336, FD-1337, FD-1338, FD-1339, FD-1340, FD-1341, FD-1342, FD-1343, FD-1344, FD-1345, FD-1346, FD-1347, FD-1348, FD-1349, FD-1350, FD-1351, FD-1352, FD-1353, FD-1354, FD-1355, FD-1356, FD-1357, FD-1358, FD-1359, FD-1360, FD-1361, FD-1362, FD-1363, FD-1364, FD-1365, FD-1366, FD-1367, FD-1368, FD-1369, FD-1370, FD-1371, FD-1372, FD-1373, FD-1374, FD-1375, FD-1376, FD-1377, FD-1378, FD-1379, FD-1380, FD-1381, FD-1382, FD-1383, FD-1384, FD-1385, FD-1386, FD-1387, FD-1388, FD-1389, FD-1390, FD-1391, FD-1392, FD-1393, FD-1394, FD-1395, FD-1396, FD-1397, FD-1398, FD-1399, FD-1400, FD-1401, FD-1402, FD-1403, FD-1404, FD-1405, FD-1406, FD-1407, FD-1408, FD-1409, FD-1410, FD-1411, FD-1412, FD-1413, FD-1414, FD-1415, FD-1416, FD-1417, FD-1418, FD-1419, FD-1420, FD-1421, FD-1422, FD-1423, FD-1424, FD-1425, FD-1426, FD-1427, FD-1428, FD-1429, FD-1430, FD-1431, FD-1432, FD-1433, FD-1434, FD-1435, FD-1436, FD-1437, FD-1438, FD-1439, FD-1440, FD-1441, FD-1442, FD-1443, FD-1444, FD-1445, FD-1446, FD-1447, FD-1448, FD-1449, FD-1450, FD-1451, FD-1452, FD-1453, FD-1454, FD-1455, FD-1456, FD-1457, FD-1458, FD-1459, FD-1460, FD-1461, FD-1462, FD-1463, FD-1464, FD-1465, FD-1466, FD-1467, FD-1468, FD-1469, FD-1470, FD-1471, FD-1472, FD-1473, FD-1474, FD-1475, FD-1476, FD-1477, FD-1478, FD-1479, FD-1480, FD-1481, FD-1482, FD-1483, FD-1484, FD-1485, FD-1486, FD-1487, FD-1488, FD-1489, FD-1490, FD-1491, FD-1492, FD-1493, FD-1494, FD-1495, FD-1496, FD-1497, FD-1498, FD-1499, FD-1500, FD-1501, FD-1502, FD-1503, FD-1504, FD-1505, FD-1506, FD-1507, FD-1508, FD-1509, FD-1510, FD-1511, FD-1512, FD-1513, FD-1514, FD-1515, FD-1516, FD-1517, FD-1518, FD-1519, FD-1520, FD-1521, FD-1522, FD-1523, FD-1524, FD-1525, FD-1526, FD-1527, FD-1528, FD-1529, FD-1530, FD-1531, FD-1532, FD-1533, FD-1534, FD-1535, FD-1536, FD-1537, FD-1538, FD-1539, FD-1540, FD-1541, FD-1542, FD-1543, FD-1544, FD-1545, FD-1546, FD-1547, FD-1548, FD-1549, FD-1550, FD-1551, FD-1552, FD-1553, FD-1554, FD-1555, FD-1556, FD-1557, FD-1558, FD-1559, FD-1560, FD-1561, FD-1562, FD-1563, FD-1564, FD-1565, FD-1566, FD-1567, FD-1568, FD-1569, FD-1570, FD-1571, FD-1572, FD-1573, FD-1574, FD-1575, FD-1576, FD-1577, FD-1578, FD-1579, FD-1580, FD-1581, FD-1582, FD-1583, FD-1584, FD-1585, FD-1586, FD-1587, FD-1588, FD-1589, FD-1590, FD-1591, FD-1592, FD-1593, FD-1594, FD-1595, FD-1596, FD-1597, FD-1598, FD-1599, FD-1600, FD-1601, FD-1602, FD-1603, FD-1604, FD-1605, FD-1606, FD-1607, FD-1608, FD-1609, FD-1610, FD-1611, FD-1612, FD-1613, FD-1614, FD-1615, FD-1616, FD-1617, FD-1618, FD-1619, FD-1620, FD-1621, FD-1622, FD-1623, FD-1624, FD-1625, FD-1626, FD-1627, FD-1628, FD-1629, FD-1630, FD-1631, FD-1632, FD-1633, FD-1634, FD-1635, FD-1636, FD-1637, FD-1638, FD-1639, FD-1640, FD-1641, FD-1642, FD-1643, FD-1644, FD-1645, FD-1646, FD-1647, FD-1648, FD-1649, FD-1650, FD-1651, FD-1652, FD-1653, FD-1654, FD-1655, FD-1656, FD-1657, FD-1658, FD-1659, FD-1660, FD-1661, FD-1662, FD-1663, FD-1664, FD-1665, FD-1666, FD-1667, FD-1668, FD-1669, FD-1670, FD-1671, FD-1672, FD-1673, FD-1674, FD-1675, FD-1676, FD-1677, FD-1678, FD-1679, FD-1680, FD-1681, FD-1682, FD-1683, FD-1684, FD-1685, FD-1686, FD-1687, FD-1688, FD-1689, FD-1690, FD-1691, FD-1692, FD-1693, FD-1694, FD-1695, FD-1696, FD-1697, FD-1698, FD-1699, FD-1700, FD-1701, FD-1702, FD-1703, FD-1704, FD-1705, FD-1706, FD-1707, FD-1708, FD-1709, FD-1710, FD-1711, FD-1712, FD-1713, FD-1714, FD-1715, FD-1716, FD-1717, FD-1718, FD-1719, FD-1720, FD-1721, FD-1722, FD-1723, FD-1724, FD-1725, FD-1726, FD-1727, FD-1728, FD-1729, FD-1730, FD-1731, FD-1732, FD-1733, FD-1734, FD-1735, FD-1736, FD-1737, FD-1738, FD-1739, FD-1740, FD-1741, FD-1742, FD-1743, FD-1744, FD-1745, FD-1746, FD-1747, FD-1748, FD-1749, FD-1750, FD-1751, FD-1752, FD-1753, FD-1754, FD-1755, FD-1756, FD-1757, FD-1758, FD-1759, FD-1760, FD-1761, FD-1762, FD-1763, FD-1764, FD-1765, FD-1766, FD-1767, FD-1768, FD-1769, FD-1770, FD-1771, FD-1772, FD-1773, FD-1774, FD-1775, FD-1776, FD-1777, FD-1778, FD-1779, FD-1780, FD-1781, FD-1782, FD-1783, FD-1784, FD-1785, FD-1786, FD-1787, FD-1788, FD-1789, FD-1790, FD-1791, FD-1792, FD-1793, FD-1794, FD-1795, FD-1796, FD-1797, FD-1798, FD-1799, FD-1800, FD-1801, FD-1802, FD-1803, FD-1804, FD-1805, FD-1806, FD-1807, FD-1808, FD-1809, FD-1810, FD-1811, FD-1812, FD-1813, FD-1814, FD-1815, FD-1816, FD-1817, FD-1818, FD-1819, FD-1820, FD-1821, FD-1822, FD-1823, FD-1824, FD-1825, FD-1826, FD-1827, FD-1828, FD-1829, FD-1830, FD-1831, FD-1832, FD-1833, FD-1834, FD-1835, FD-1836, FD-1837, FD-1838, FD-1839, FD-1840, FD-1841, FD-1842, FD-1843, FD-1844, FD-1845, FD-1846, FD-1847, FD-1848, FD-1849, FD-1850, FD-1851, FD-1852, FD-1853, FD-1854, FD-1855, FD-1856, FD-1857, FD-1858, FD-1859, FD-1860, FD-1861, FD-1862, FD-1863, FD-1864, FD-1865, FD-1866, FD-1867, FD-1868, FD-1869, FD-1870, FD-1871, FD-1872, FD-1873, FD-1874, FD-1875, FD-1876, FD-1877, FD-1878, FD-1879, FD-1880, FD-1881, FD-1882, FD-1883, FD-1884, FD-1885, FD-1886, FD-1887, FD-1888, FD-1889, FD-1890, FD-1891, FD-1892, FD-1893, FD-1894, FD-1895, FD-1896, FD-1897, FD-1898, FD-1899, FD-1900, FD-1901, FD-1902, FD-1903, FD-1904, FD-1905, FD-1906, FD-1907, FD-1908, FD-1909, FD-1910, FD-1911, FD-1912, FD-1913, FD-1914, FD-1915, FD-1916, FD-1917, FD-1918, FD-1919, FD-1920, FD-1921, FD-1922, FD-1923, FD-1924, FD-1925, FD-1926, FD-1927, FD-1928, FD-1929, FD-1930, FD-1931, FD-1932, FD-1933, FD-1934, FD-1935, FD-1936, FD-1937, FD-1938, FD-1939, FD-1940, FD-1941, FD-1942, FD-1943, FD-1944, FD-1945, FD-1946, FD-1947, FD-1948, FD-1949, FD-1950, FD-1951, FD-1952, FD-1953, FD-1954, FD-1955, FD-1956, FD-1957, FD-1958, FD-1959, FD-1960, FD-1961, FD-1962, FD-1963, FD-1964, FD-1965, FD-1966, FD-1967, FD-1968, FD-1969, FD-1970, FD-1971, FD-1972, FD-1973, FD-1974, FD-1975, FD-1976, FD-1977, FD-1978, FD-1979, FD-1980, FD-1981, FD-1982, FD-1983, FD-1984, FD-1985, FD-1986, FD-1987, FD-1988, FD-1989, FD-1990, FD-1991, FD-1992, FD-1993, FD-1994, FD-1995, FD-1996, FD-1997, FD-1998, FD-1999, FD-2000, FD-2001, FD-2002, FD-2003, FD-2004, FD-2005, FD-2006, FD-



Severe

MAXIMUM STRENGTH

Nighttime

Cold/Flu Relief

Compare to the active ingredients in
Vicks[®] NyQuil[®] Severe Cold & Flu[®]

Severe

NDC 51013-110-04

CVS 11-41

THIS PRODUCT IS PACKAGED IN A CHILD RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF INSTRUCTIONS ARE PRESENT.

STORE AT ROOM TEMPERATURE.
AVOID EXCESSIVE HEAT.
KEEP OUT OF REACH OF CHILDREN.
WARNING: Avoid product ingestion.



MAXIMUM STRENGTH

Nighttime

Cold/Flu Relief

ACETAMINOPHEN - Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
DOXYLAMINE SUCCINATE - Antihistamine
PHENYLEPHRINE HCl - Nasal decongestant

Relieves:
 Headache; Fever; Coughing; Sneezing;
 Runny nose; Nasal congestion.

24 SOFTGELS

Actual Size

LOT NO. COATING
EXP. DATE FREE AREA

31

This product is not manufactured or distributed by Pkce, Inc. (owner of the registered trademark "NyQuil Severe" & its variations or comments) 1-888-275-8100

Active ingredients (in each softgel):
 Acetaminophen 325 mg, Dextromethorphan HBr 10 mg, Doxylamine Succinate 2 mg, Phenylephrine HCl 5 mg

Other information:
 • Store at room temperature and avoid excessive heat.
 • Keep out of reach of children.
 • Use only if instructions are present.

Drug Facts (continued):
 • When you go to bed at nighttime and take this product, take it as directed to ensure correct dosing.

LOT NO. COATING
EXP. DATE FREE AREA

COATING FREE AREA

Drug Facts

Active ingredients (in each softgel):
 Acetaminophen 325 mg, Dextromethorphan HBr 10 mg, Doxylamine Succinate 2 mg, Phenylephrine HCl 5 mg

Other information:
 • Store at room temperature and avoid excessive heat.
 • Keep out of reach of children.
 • Use only if instructions are present.

Drug Facts (continued):
 • When you go to bed at nighttime and take this product, take it as directed to ensure correct dosing.

LOT NO. COATING
EXP. DATE FREE AREA

COATING FREE AREA

Drug Facts

Active ingredients (in each softgel):
 Acetaminophen 325 mg, Dextromethorphan HBr 10 mg, Doxylamine Succinate 2 mg, Phenylephrine HCl 5 mg

Other information:
 • Store at room temperature and avoid excessive heat.
 • Keep out of reach of children.
 • Use only if instructions are present.

Drug Facts (continued):
 • When you go to bed at nighttime and take this product, take it as directed to ensure correct dosing.

LOT NO. COATING
EXP. DATE FREE AREA

COATING FREE AREA


Drug Facts

Active ingredients (in each softgel):
 Acetaminophen 325 mg, Dextromethorphan HBr 10 mg, Doxylamine Succinate 2 mg, Phenylephrine HCl 5 mg

Other information:
 • Store at room temperature and avoid excessive heat.
 • Keep out of reach of children.
 • Use only if instructions are present.

Drug Facts (continued):
 • When you go to bed at nighttime and take this product, take it as directed to ensure correct dosing.

LOT NO. COATING
EXP. DATE FREE AREA



Nighttime Severe Cold & Cough

Drug Facts	
Active ingredients (In each packet)	Purposes
Acetaminophen 500 mg	Pain relief/fever reducer
Diphenhydramine	Sleep aid
hydrochloride 25 mg	Antihistamine/Cough Suppressant
Phenylephrine hydrochloride 10 mg	Nasal Decongestant
Uses	
<ul style="list-style-type: none"> ■ temporarily relieves these symptoms due to a cold: ■ minor aches and pains ■ minor sore throat pain ■ headache ■ nasal and sinus congestion ■ runny nose ■ sneezing ■ itchy nose or throat ■ itchy, watery eyes due to hay fever ■ cough due to minor throat and bronchial irritation ■ temporarily reduces fever 	

Warnings
Liver warning: This p

Severe liver damage may occur if you take **more than 4,000 mg of acetaminophen** in 24 hours. **Do not take** **more than 4,000 mg of acetaminophen** in 24 hours. **Do not take** **more than 3,000 mg of acetaminophen** in 24 hours if you are taking **other drugs containing acetaminophen** or **3 or more alcoholic drinks every day while using this product**.

Warning: **Alert:** acetaminophen may cause severe skin reactions. Symptoms may include **skin redness** or **itching**.

Caution: **Alert:** If a skin reaction occurs, stop use and seek medical help right away.

See Warnings: If throat is sore, severe, persists for more than 7 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor immediately.

Do not use: **Do not use** in a child under 17 years of age. **Do not use** if you are allergic to acetaminophen **or** with any other drug containing acetaminophen (prescription and nonprescription). If you are taking any other drug, ask your doctor if you can take the drug **or** pharmacist **or** with any other product containing dicyclanil/diphenhydramine, even one used on the skin.

Do not use: **Do not use** if you are taking or have taken an MAO inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after you have taken an MAOI. **Do not use** if you are taking your prescription drug contains a MAOI, ask your doctor or pharmacist if you can take this product.

Ask a doctor before you stop using this product.

Do not use:

- Do not use** if you have liver disease
- Do not use** if you have heart disease
- Do not use** if you have high blood pressure
- Do not use** if you have thyroid disease
- Do not use** if you have diabetes
- Do not use** if you have glucose in your urine

Multi-Symptom Severe Cold

Drug Facts	
Active Ingredients (in each packet)	Purposes
Acetaminophen 500 mg	Pain relief/fever reduction
Dextromethorphan	
Hydrocodone 20 mg	Cough Suppressant
Phenylephrine hydrochloride 10 mg	Nasal Decongestant

Uses
Temporarily relieves these symptoms due to a cold:
• minor aches and pains
• minor sore throat pain
• headache
• nasal and sinus congestion
• cough
• minor throat and bronchial irritation
Temporarily reduces fever

Warnings
Do not take if you are already taking another medicine that contains acetaminophen or hydrocodone.
Do not take if you are taking any medicine for depression or anxiety.
Do not take if you are taking any medicine for high blood pressure.
Do not take if you are taking any medicine for heart disease.
Do not take if you are taking any medicine for diabetes.
Do not take if you are taking any medicine for thyroid disease.
Do not take if you are taking any medicine for kidney disease.
Do not take if you are taking any medicine for liver disease.
Do not take if you are taking any medicine for stomach or intestinal disease.
Do not take if you are taking any medicine for urinary tract disease.
Do not take if you are taking any medicine for blood clotting.
Do not take if you are taking any medicine for blood sugar.
Do not take if you are taking any medicine for blood pressure.
Do not take if you are taking any medicine for heart rate.
Do not take if you are taking any medicine for cholesterol.
Do not take if you are taking any medicine for triglycerides.
Do not take if you are taking any medicine for calcium.
Do not take if you are taking any medicine for iron.
Do not take if you are taking any medicine for zinc.
Do not take if you are taking any medicine for vitamin D.
Do not take if you are taking any medicine for vitamin E.
Do not take if you are taking any medicine for vitamin K.
Do not take if you are taking any medicine for vitamin B.
Do not take if you are taking any medicine for vitamin C.
Do not take if you are taking any medicine for vitamin A.
Do not take if you are taking any medicine for vitamin D.
Do not take if you are taking any medicine for vitamin E.
Do not take if you are taking any medicine for vitamin K.
Do not take if you are taking any medicine for vitamin B.
Do not take if you are taking any medicine for vitamin C.
Do not take if you are taking any medicine for vitamin A.

Liver warning:
Severe liver damage
4,000 mg of acetaminophen

drugs containing acetaminophen. **■** If you're taking more than one dose every 4 hours while using this product, you may be taking too much acetaminophen, which can cause severe liver damage. Symptoms may include:

- loss of appetite
- nausea
- stomach pain
- dark urine
- yellowing of the skin or the whites of the eyes

If you notice reaction, stop use and seek medical help.

Take special warning: If you're taking it seven, possibly for more than 7 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor.

Somebody else's medicine:

- **Do not use**
- if you're in a child under 12 years of age
- if you're allergic to acetaminophen
- if you're taking other drugs containing acetaminophen (prescription and nonprescription), if you're not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you're taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, Parkinson's, or emotional conditions, or Parkinson's)
- if you're taking any medicine that says "avoid alcohol" or "do not drink alcohol" while taking the drug
- if you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before using it you have

- liver or kidney disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

Multi-Symptom Severe Cold (Continued)

Drug Facts (continued)

• cough that occurs with too much phlegm (mucus)
 • ask a doctor if there is chronic such as occurs with smoking, asthma or emphysemas.

Ask a doctor or pharmacist before use if you are: taking the blood thinning drug warfarin
 • taking any other medicine
 • pregnant or nursing
 • taking or about to take alcohol or grapefruit juice

Stop use and ask a doctor if:
 • new rashes, dizziness, or drowsiness occurs
 • new or worse heartburn or indigestion occurs
 • redness or swelling is present
 • symptoms do not get better or worsen
 • you feel, cough or need chest congestion gets worse or lasts more than 7 days
 • cough comes back or occurs, rash or headache that lasts.

Keep out of reach of children. Stop use and ask a doctor if:
 • pregnant or breast-feeding, ask a health care professional before use.
 • keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (See back of package for more information.)
 • ask a doctor if you are taking or about to take any medicine or if you are taking or about to take any medicine or if you are taking or about to take any medicine.

Directions
 ■ do not use more than d

[illegible]**VALUE PACK**



Compare to the active ingredients in Therma® Multi-Symptom Severe Cold & Fluallergies, Severe Cold & Cough, and Severe Allergies.

Severe Cold & Cough

NDC: 527.75-758

Multi-Symptom

Severe Cold

ACETAMINOPHEN
Pseudoephedrine HCl
DEXTROMETHORPHAN HBr
Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:

- Nasal congestion
- Sore throat pain
- Cough
- Headache
- Body aches
- Fever




6 PACKETS

Nighttime

Severe Cold & Cough

ACETAMINOPHEN
Diphenhydramine HCl
DIPHENHYDRAMINE HCl
Antihistamine, Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:

- Nasal congestion
- Sore throat pain
- Cough
- Headache
- Body aches
- Fever
- Rummy nose
- Sneezing




6 PACKETS

Nighttime Severe Cold & Cough (Continued)

Drug Facts (continued)

- Trouble arising due to an enlarged prostate gland
- Breathing problem such as emphysema, asthma or chronic bronchitis
- Drowsiness or dizziness
- Drowsiness that occurs with alcohol, other drugs or grapefruit juice
- Drowsiness that leads to dizziness such as smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are taking:

- sedatives or tranquilizers
- the blood thinning drug warfarin

When using this product:

- **Do not exceed recommended dose.** Mild alcoholic drinks **may** mask drowsiness or other side effects.
- Alcohol, sedatives and tranquilizers may increase drowsiness. **Be careful** when driving a motor vehicle or operating machinery. **Excess** use may cause harm, especially in children.

Stop use and ask a doctor if:

- nervousness, dizziness, or drowsiness occurs
- new fever or rash or lasts more than 7 days
- drowsiness or swelling is present
- symptoms do not get better or worsen
- pain, cough or chest congestion gets worse or lasts more than 7 days

These could be signs of

Directions

We do not eat what's inside

Give 1 packet 3 times each appropriate period, but do not exceed 6 packets in 24 hours unless advised by a doctor

Age	One packet	Three
Adults and children 15 years of age and older	One packet	Three
Children under 15 years of age	Do not use	Do not use

Use the contents of one packet into 8 oz. of water, stir; spit white hot. Consume entire drink. Do not eat or drink anything else for 1 hour after use. Do not use more than 6 packets in 24 hours. If you are taking any other medicine, ask your doctor if you should stop taking it. If you are taking a microwave, add contents of one packet to 8 oz. of cool water; mix briefly before heating. Do not use in a microwave oven.

Other information

- Each packet contains: potassium 10 mEq
- Contains 100 mg of sodium
- Protected from excessive heat and moisture

Inactive ingredients are: croscarmellose sodium, PEG 400, yellow #6, flavor, maltodextrin, potassium chloride, and sodium citrate

Questions or comments ? 800-827-2718

© 2005 Pharmacia Corporation. All rights reserved. Manufactured by Pharmacia Corporation, Kalamazoo, MI. All rights reserved. All trademarks are the property of their respective owners. All other trademarks are the property of their respective owners. All other trademarks are the property of their respective owners.

Lot:

Exp

54812CVSFC



Distributed by:
CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 028
© 2017 CVS/pharma
CVS.com®
1-800-SHOP CVS
V-12431



**TAMPER EVIDENT INNER UNIT: DO NOT USE IF
SEALED PACKET IS TORN OR BROKEN.**



Compare to the active ingredients in Alka-Seltzer PLUS® Maximum Strength Severe Sinus Congestion & Cough Day & Night Liquid Gels*

Sinus Congestion & Cough

Daytime Non-Drowsy

Sinus Congestion & Cough Relief

ACETAMINOPHEN 325 mg
Pain reliever/Fever reducer
DEXTROMETHORPHAN HBr 10 mg
Cough suppressant
PHENYLEPHRINE HCl 5 mg
Nasal decongestant

Relieves:

Nasal & sinus congestion;
Sinus pressure; Headache &
body ache; Cough



Actual Size

12 SOFTGELS

Nighttime

Sinus Congestion & Cough Relief

ACETAMINOPHEN 325 mg
Pain reliever/Fever reducer
DEXTROMETHORPHAN HBr 10 mg
Cough suppressant
DOXYLAMINE SUCCINATE 6.25 mg
Antihistamine
PHENYLEPHRINE HCl 5 mg
Nasal decongestant

Relieves:

Sinus congestion &
pressure; Headache &
pain; Runny nose &
sneezing; Cough



Actual Size

8 SOFTGELS 20 TOTAL

CVS Health.

DAYTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
GUAFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant
6 DAY PACKETS

NIGHTTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
DOXYLAMINE SUCCLATE
Antihistamine
PHENYLEPHRINE HCl
Nasal decongestant
6 NIGHT PACKETS
TOTAL 12 PACKETS

CVS Health.

Compare to the active ingredients in
Alka-Seltzer PLUS® Severe Cold & Flu Day and Night

DAYTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
GUAFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant
6 DAY PACKETS

NIGHTTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
DOXYLAMINE SUCCLATE
Antihistamine
PHENYLEPHRINE HCl
Nasal decongestant
6 NIGHT PACKETS
TOTAL 12 PACKETS

Hot Lemon Flavor
Naturally & Artificially Flavored

Hot Lemon Flavor
Naturally & Artificially Flavored

6 DAY PACKETS

6 NIGHT PACKETS
TOTAL 12 PACKETS

Lot: _____

Exp: _____

#009420

50428 584 17

Compare to the active ingredients in
Alka-Seltzer PLUS® Severe Cold & Flu Day and Night

DAYTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
GUAFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant
6 DAY PACKETS

Compare to the active ingredients in
Alka-Seltzer PLUS® Severe Cold & Flu Day and Night

NIGHTTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
DOXYLAMINE SUCCLATE
Antihistamine
PHENYLEPHRINE HCl
Nasal decongestant
6 NIGHT PACKETS
TOTAL 12 PACKETS

Compare to the active ingredients in
Alka-Seltzer PLUS® Severe Cold & Flu Day and Night

DAYTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
GUAFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant
6 DAY PACKETS

Compare to the active ingredients in
Alka-Seltzer PLUS® Severe Cold & Flu Day and Night

NIGHTTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
DOXYLAMINE SUCCLATE
Antihistamine
PHENYLEPHRINE HCl
Nasal decongestant
6 NIGHT PACKETS
TOTAL 12 PACKETS

Compare to the active ingredients in
Alka-Seltzer PLUS® Severe Cold & Flu Day and Night

DAYTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
GUAFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant
6 DAY PACKETS

Compare to the active ingredients in
Alka-Seltzer PLUS® Severe Cold & Flu Day and Night

NIGHTTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
DOXYLAMINE SUCCLATE
Antihistamine
PHENYLEPHRINE HCl
Nasal decongestant
6 NIGHT PACKETS
TOTAL 12 PACKETS

Compare to the active ingredients in
Alka-Seltzer PLUS® Severe Cold & Flu Day and Night

DAYTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
GUAFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant
6 DAY PACKETS

Compare to the active ingredients in
Alka-Seltzer PLUS® Severe Cold & Flu Day and Night

NIGHTTIME
Severe Cold & Flu
ACETAMINOPHEN
Pain reliever/ Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
DOXYLAMINE SUCCLATE
Antihistamine
PHENYLEPHRINE HCl
Nasal decongestant
6 NIGHT PACKETS
TOTAL 12 PACKETS





Compare to the active ingredients in
Vicks® NyQuil® Severe Cold & Flu*

MAXIMUM STRENGTH

Severe Nighttime

Cold/Flu Relief

ACETAMINOPHEN - Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr - Cough suppressant

DOXYLAMINE SUCCINATE - Antihistamine

PHENYLEPHRINE HCl - Nasal decongestant

Relieves:

Headache; Fever; Coughing;
Sneezing; Runny nose;
Nasal congestion



Actual Size

24 SOFTGELS



Drug Facts

Active ingredients (in each packet) **Purpose**
 Acetaminophen 650 mg.....Pain reliever/Fever reducer
 Diphenhydramine hydrochloride 25 mg.....Antihistamine/Cough suppressant
 Phenylephrine hydrochloride 10 mg.....Nasal decongestant

Uses

- Temporarily relieves:
 - minor aches and pains • headache • minor sore throat pain
 - nasal and sinus congestion • cough due to minor throat and bronchial irritation • runny nose • sneezing • itchy, watery eyes due to hay fever • itchy nose and throat
- Temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 6 packets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription and nonprescription).
- Ask a doctor or pharmacist before using with other drugs if you are not sure.
- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure • thyroid disease
- glaucoma • diabetes • trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma, or chronic bronchitis
- cough that occurs with too much phlegm (mucus) • cough that lasts or is chronic such as occurs with smoking, asthma or emphysema.

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers • taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks • marked drowsiness may occur • alcohol, sedatives and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children.

Stop use and ask a doctor if

- fever, rash, or skin redness occurs • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur
- symptoms do not get better or worsen • pain, cough or nasal congestion gets worse or lasts more than 7 days • cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- take every 4 hours; not to exceed 6 packets in 24 hours or as directed by a doctor.
- adults and children 12 years of age and over: dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years of age: consult a doctor.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 7 mg, sodium 48 mg
- phenylephrine: contains phenylalanine 13 mg per packet
- store at controlled room temperature 20-25°C (68-77°F). Protect from excessive heat and moisture.

Inactive ingredients:

Acetosulfame K, Aspartame, Caramel Color, Citric Acid, D&C Yellow 10, Maltodextrin, Natural Flavors, Sodium Citrate, Starch and Sugar.

DO NOT USE IF SEALED PACKET IS TORN OR BROKEN.

Distributed by: CVS Pharmacy, Inc.
 One CVS Drive, Woonsocket, RI 02895
 © 2013 CVS Pharmacy
 www.cvs.com 1-800-shop-CVS
 Made in Canada
 V-16833

CVS Quality
 Money Back Guarantee

SEVERE COUGH & COLD NIGHTTIME

CVS
pharmacy

NDC 59779-221-06

NEW!

SEVERE COUGH & COLD NIGHTTIME

ACETAMINOPHEN – PAIN RELIEVER / FEVER REDUCER
DIPHENHYDRAMINE HCl – ANTIHISTAMINE / COUGH SUPPRESSANT
PHENYLEPHRINE HCl – NASAL DECONGESTANT

- Nasal congestion
- Sore throat pain
- Cough
- Sneezing
- Runny nose
- Headache
- Body ache
- Fever

Honey Lemon Infused
 with Chamomile &
 White Tea flavors

6 PACKETS

SEE NEW WARNINGS INFORMATION

CVS
pharmacy

SEVERE COUGH & COLD NIGHTTIME

CVS/pharmacy™ Severe Cough & Cold Nighttime provides complete relief of your cough and severe cold symptoms. This soothing hot liquid formula is alcohol free and tastes great.

For daytime relief, try

C-3109-30508-6D

<p>Living Facts Call it a Day!</p> <p>Active symptoms Passive symptoms</p> <p>Active symptoms are those that are causing the disease. Passive symptoms are those that are caused by the disease.</p> <p>Example: A person with a cold has a fever. The fever is a passive symptom because it is caused by the cold.</p>	<p>Living Facts Barry Corcoran's Deep Breath</p> <p>Active symptoms Passive symptoms</p> <p>Active symptoms are those that are causing the disease. Passive symptoms are those that are caused by the disease.</p> <p>Example: A person with a cold has a fever. The fever is a passive symptom because it is caused by the cold.</p>	<p>Living Facts Barry Corcoran's Deep Breath</p> <p>Active symptoms Passive symptoms</p> <p>Active symptoms are those that are causing the disease. Passive symptoms are those that are caused by the disease.</p> <p>Example: A person with a cold has a fever. The fever is a passive symptom because it is caused by the cold.</p>
<p>Living Facts Barry Corcoran's Deep Breath</p> <p>Active symptoms Passive symptoms</p> <p>Active symptoms are those that are causing the disease. Passive symptoms are those that are caused by the disease.</p> <p>Example: A person with a cold has a fever. The fever is a passive symptom because it is caused by the cold.</p>	<p>Living Facts Barry Corcoran's Deep Breath</p> <p>Active symptoms Passive symptoms</p> <p>Active symptoms are those that are causing the disease. Passive symptoms are those that are caused by the disease.</p> <p>Example: A person with a cold has a fever. The fever is a passive symptom because it is caused by the cold.</p>	<p>Living Facts Barry Corcoran's Deep Breath</p> <p>Active symptoms Passive symptoms</p> <p>Active symptoms are those that are causing the disease. Passive symptoms are those that are caused by the disease.</p> <p>Example: A person with a cold has a fever. The fever is a passive symptom because it is caused by the cold.</p>

2711

DAY & NIGHT COMBO PACK

CVS Health.

MAXIMUM STRENGTH Daytime Severe Cold

Acetaminophen
Pain reliever/Fever reducer
Dextromethorphan HBr
Cough suppressant
Guaifenesin, Expectoant
Phenylephrine HCl
Nasal decongestant

MULTI-SYMPTOM

Headache + Body Pain
Fever + Sore Throat
Cough + Chest Congestion
Nasal Congestion

FOR AGES 12+

Actual Size

20 DAYTIME CAPLETS
30 TOTAL CAPLETS

MAXIMUM STRENGTH Nighttime Cold & Flu

Acetaminophen
Pain reliever/Fever reducer
Diphenhydramine HCl
Antihistamine
Phenylephrine HCl
Nasal decongestant

MULTI-SYMPTOM

Headache + Body Pain
Fever + Sore Throat
Nasal Congestion
Sneezing & Runny Nose

FOR AGES 12+

Actual Size

10 NIGHTTIME CAPLETS

CVS Health.

MAXIMUM STRENGTH Daytime Severe Cold

Acetaminophen,
Dextromethorphan HBr,
Guaifenesin, Phenylephrine HCl

MAXIMUM STRENGTH Nighttime Cold & Flu

Acetaminophen,
Diphenhydramine HCl,
Phenylephrine HCl

No print/No varnish area
Lot no/Exp date

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Do not take the
Daytime and Nighttime
caplets at the same time.

ADHESIVE AREA

Drug Facts (continued)

■ if you have ever had an allergic reaction to this product or any of its ingredients
■ with any other product containing diphenhydramine, even one used on skin (**Nighttime only**)

Ask a doctor before use if you have

- heart disease ■ thyroid disease ■ diabetes
- liver disease ■ high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis (**Nighttime only**) ■ glaucoma (**Nighttime only**)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**Nighttime only**)

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children (**Nighttime only**)
- marked drowsiness may occur (**Nighttime only**)

TYPE SAFETY

ADHESIVE AREA

Drug Facts (continued)

■ avoid alcoholic beverages (**Nighttime only**)
■ alcohol, sedatives, and tranquilizers may increase drowsiness (**Nighttime only**)
■ use caution when driving a motor vehicle or operating machinery (**Nighttime only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Do not take DAYTIME and NIGHTTIME products at the same time.

ADHESIVE AREA

Drug Facts (continued)

Directions ■ do not take more than directed
■ do not take more than 12 caplets in any 24-hour period
■ adults and children 12 years and over: take 2 caplets every 4 hours
■ children under 12 years: do not use

Other information
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number

Inactive ingredients (Daytime only) corn starch, croscopolone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

ADHESIVE AREA

Drug Facts (continued)

Inactive ingredients (Nighttime only) corn starch, croscopolone, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide

Questions or comments? 1-800-426-9391

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-19849

100% money back guaranteed.
CVS.com/returnpolicy

50044
ORG102161769401
R-0231-617694-01H-DF

Allergy Plus Congestion Relief

DIPHENHYDRAMINE HCl - Antihistamine
PHENYLEPHRINE HCl - Nasal decongestant

Relief of:

- Sneezing
- Runny nose
- Sinus congestion
- Sinus pressure

Tablets

CVS Health

Compare to the active ingredients in Benadryl® Allergy Plus Congestion Relief

Tablets

CVS Health

Allergy Plus Congestion Relief

DIPHENHYDRAMINE HCl - Antihistamine
PHENYLEPHRINE HCl - Nasal decongestant

B-0231-485-08-H
ORG122048508

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredients (in each tablet)

Diphenhydramine HCl 25 mg Antihistamine
Phenylephrine HCl 10 mg Nasal decongestant

Purpose

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - nasal congestion
 - itching of the nose or throat
 - itchy, watery eyes
- temporarily relieves these symptoms due to the common cold:
 - nasal congestion
 - runny nose
 - sneezing
- temporarily relieves sinus congestion and pressure

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- diabetes
- thyroid disease
- heart disease
- glaucoma
- high blood pressure
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

Drug Facts (continued)

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- symptoms do not improve within 7 days or are accompanied by fever
- nervousness, dizziness, or sleeplessness occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- adults and children 12 years and over
 - take 1 tablet every 4 hours
 - do not take more than 6 tablets in 24 hours
- children under 12 years: do not use

Other information

- each tablet contains: calcium 25 mg
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, hypromellose, magnesium stearate,

Drug Facts (continued)

microcrystalline cellulose, polydextrose, polyethylene glycol, silicon dioxide, stearic acid, titanium dioxide, triacetin

Questions or comments? 1-800-426-9391

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Benadryl® Allergy Plus Congestion.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02896
© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-18615

100% money back guaranteed.
CVS.com/returnpolicy



CVS Health

Tablets


Allergy Plus Congestion Relief

DIPHENHYDRAMINE HCl - Antihistamine
PHENYLEPHRINE HCl - Nasal decongestant

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No Print/No Varnish
Lot and Expiration No.

#197895
FP0 100%
UPC# 050428455623
XXXXXX XXXXXX



COMBO PACK

Drug Facts DAYTIME

Active Ingredients (in each 15 mL)

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Phenylephrine HCl 5 mg Nasal decongestant

Uses ■ Temporarily relieves common cold and flu symptoms:
■ sore throat ■ nasal congestion ■ minor aches and pains
■ cough due to minor throat and bronchial irritation ■ headache
■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if:
■ you take more than 4,000 mg of acetaminophen in 24 hours
■ you take more than 5 doses in 24 hours
■ you take with other drugs containing acetaminophen
■ you take 5 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ hives ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (paracetamol or paracetamol). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if the user has
■ difficulty in urination due to enlargement of the prostate gland
■ cough that occurs with too much phlegm (mucus)
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ heart disease ■ diabetes ■ thyroid disease ■ liver disease ■ high blood pressure
Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin.

Drug Facts (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if
■ hives, rash, or skin redness occur
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present ■ new symptoms occur
■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (4 days for 7 days old)
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ **do not take more than directed**
■ do not exceed 4 doses per 24 hours
■ mL = milliliter ■ **Must use the dose as provided**

adults and children 12 years and older	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

Other information
■ each 15 mL contains: sodium 13 mg
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ use by expiration date on package

Inactive ingredients hydroxy acid, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, drug-free sodium saccharin, sodium saccharin, sorbitol, sucralose.

Questions or comments? 1-800-426-9391

Non-Drowsy Daytime

COLD & FLU

ACETAMINOPHEN
Pain reliever, Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
PHENYLEPHRINE HCl
Nasal decongestant

Relieves: Headache, Fever, Sore throat, Minor aches & pains, Nasal congestion, Sinus pressure, Cough

Cherry Flavor
No added alcohol
2 - 12 FL OZ (355 mL) BOTTLES

Multi-Symptom Nighttime

COLD & FLU

ACETAMINOPHEN
Pain reliever, Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
DOXYLAMINE SUCCINATE
Antihistamine

Relieves: Headache, Fever, Sore throat, Minor aches & pains, Sneezing, Runny nose, Cough

Cherry Flavor
No added alcohol
TOTAL 24 FL OZ (710 mL)

Drug Facts NIGHTTIME

Active Ingredients (in each 30 mL)

Acetaminophen 650 mg Pain reliever/fever reducer
Dextromethorphan HBr 20 mg Cough suppressant
Doxylamine succinate 12.5 mg Antihistamine

Uses ■ Temporarily relieves common cold and flu symptoms:
■ headache ■ minor aches and pains
■ sore throat ■ runny nose and sneezing
■ cough due to minor throat and bronchial irritation
■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 5 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ hives ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (paracetamol or paracetamol). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if you have
■ difficulty in urination due to enlargement of the prostate gland
■ cough that occurs with too much phlegm (mucus)
■ a breathing problem or chronic cough that lasts or so occurs with mucus, asthma, chronic bronchitis, or emphysema ■ diabetes
Ask a doctor or pharmacist before use if you are
■ taking the blood thinning drug warfarin
■ taking sedatives or tranquilizers

When using this product
■ do not take more than directed
■ avoid alcoholic beverages ■ marked drowsiness may occur
■ do not drink, eat, or take any other medicine while using this product
■ use caution when driving a motor vehicle or operating machinery
Sing use and ask a doctor if
■ you or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present ■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ **do not take more than directed**
■ do not exceed 4 doses per 24 hours
■ mL = milliliter ■ **Only use the dose as provided**
■ adults and children 12 years and older: 30 mL every 6 hours
■ children under 12 years: do not use

Other information ■ each 30 mL contains: sodium 13 mg
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ use by expiration date on package

Inactive ingredients hydroxy acid, FD&C blue #1, FD&C red #40, flavor, glycerin, high fructose corn syrup, polyethylene glycol, 400, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium citrate dihydrate, sodium saccharin, sucralose.

Questions or comments? 1-800-426-9391

PARENTS:
www.diphenhydramine.org

"This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks, DayQuil, Cold & Flu, NyQuil, Cold & Flu."

#180153

FFD 80%
UPCH 060426295180
XXXXXX
W42031-01213-427W

NO PRINT / NO VARNISH AREA
Lot # and Exp. Info



Compare to the
active ingredients
in Vicks® DayQuil®
Severe Honey Flavor*

NDC 51316-613-40

Daytime Severe

ACETAMINOPHEN

Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr

Cough suppressant

GUAIFENESIN Expectorant

PHENYLEPHRINE HCl

Nasal decongestant

Relieves:

Headache, Fever, Sore throat, Minor
aches & pains, Chest congestion,
Nasal & Sinus congestion, Sinus
pressure, Cough

Thins & loosens mucus

COLD & FLU



**Honey
Flavor**

ANTIHISTAMINE FREE

12 FL OZ (355 mL)

: 51040 17 F1

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive, Woonsocket,
RI 02895 © 2023 CVS/pharmacy
CVS.com* 1-800-SHOP CVS V-16430

100% money back
guaranteed.
CVS.com/returnpolicy

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

DO NOT USE IF PRINTED NECKBAND
IS BROKEN OR MISSING.

Empty &
Replace Cap



how2recycle.info

#350947

Drug Facts

Active ingredients (in each 15 mL)

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/ fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses temporarily relieves common cold/flu
symptoms: ■ sinus congestion and pressure

■ nasal congestion ■ minor aches and
pains ■ headache ■ sore throat
■ fever ■ cough due to minor throat and bronchial irritation
■ reduces swelling of nasal passages ■ promotes nasal
and/or sinus drainage ■ temporarily restores freer breathing
through the nose ■ helps loosen phlegm (mucus) and thin
bronchial secretions to rid the bronchial passageways of
bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen.
Severe liver damage may occur if ■ adult takes more than
4,000 mg of acetaminophen in 24 hours ■ child takes
more than 5 doses in 24 hours ■ taken with other drugs
containing acetaminophen ■ adult has 3 or more
alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin
reactions. Symptoms may include: ■ skin reddening

■ blisters ■ rash
If a skin reaction occurs,
stop use and seek medical
help right away.

PEEL BACK AT
CORNER FOR MORE
INFORMATION

: 51040 17 B1

Drug Facts (continued)

Sore throat warning: If sore throat
is severe, persists for more than 2 days,
is accompanied or followed by fever,
headache, rash, nausea, or vomiting,
consult a doctor promptly.

Do not use ■ with any other drug
containing acetaminophen (prescription
or nonprescription). If you are not sure
whether a drug contains acetaminophen,
ask a doctor or pharmacist.

■ if you are now taking a prescription
monoamine oxidase inhibitor (MAOI)
(certain drugs for depression,
psychiatric, or emotional conditions, or
Parkinson's disease), or for 2 weeks
after stopping the MAOI drug. If you do
not know if your prescription drug
contains an MAOI, ask a doctor or
pharmacist before taking this product.
■ if you have ever had an allergic
reaction to this product or any of its
ingredients

Ask a doctor before use if you have

Drug Facts (continued)

Stop use and ask a doctor if ■ you get
nervous, dizzy or sleepless ■ pain, nasal
congestion, or cough gets worse or lasts more than
5 days (children) or 7 days (adults) ■ fever gets
worse or lasts more than 3 days ■ redness or
swelling is present ■ new symptoms occur
■ cough comes back or occurs with rash or headache
that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health
professional before use.


Keep out of reach of children. Overdose warning:
In case of overdose, get medical help or contact a
Poison Control Center right away (1-800-222-1222).
Quick medical attention is critical for adults as well
as for children even if you do not notice any signs or
symptoms.

Directions ■ take only as directed - see
Overdose warning ■ only use the dose cup
provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs

ADHESIVE AREA
• NO VARNISH • NO TYPE

ADHESIVE AREA
• NO VARNISH • NO TYPE



Severe

Ingredient	Daytime	Nighttime
Acetaminophen	85%	90%
Phenylephrine HCl	75%	80%

Cold & Flu Relief

Ingredient	Daytime	Nighttime
Acetaminophen	75%	80%
Phenylephrine HCl	65%	70%

CVS Health.

[illegible]

B-0231-640677-22H
REV0722F64067722

← Lot and Exp will get printed on the line here

CVS Health. Compare to the active ingredients in Vicks® NyQuil® Severe Honey Flavor*
NDC 51316-179-40

Nighttime Severe

ACETAMINOPHEN
Pain reliever; Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
DOXYLAMINE SUCCINATE
Antihistamine
PHENYLEPHRINE HCl
Nasal decongestant

Relieves:
Fever, Sore throat, Minor aches & pains, Nasal congestion, Sinus pressure, Sneezing, Runny nose, Cough

COLD & FLU

Honey Flavor

12 FL OZ (355 mL)

: 8LS4D 17 F1

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive, Woonsocket,
RI 02895 © 2023 CVS/pharmacy
CVS.com* 1-800-SHOP CVS V-16430
100% money back guaranteed.
CVS.com/returnpolicy

PARENTS: **DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.**
Learn about teen medicine abuse
www.StopMedicineAbuse.org

#545974

0 50428 56697 8

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Doxylamine Succinate 6.25 mg.....	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ cough to help you sleep ■ minor aches and pains ■ headache ■ sore throat ■ fever ■ runny nose and sneezing ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage

Warnings
Liver warnings: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warnings: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

PEEL BACK AT CORNER FOR MORE INFORMATION

: 8LS4D 17 B1

Drug Facts (continued)

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients ■ to make a child sleepy

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ glaucoma ■ thyroid disease ■ diabetes ■ cough that occurs with too much

ESIVE AREA NO VARNISH • NO TYPE

Drug Facts (continued)

■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

ADHESIVE AREA NO VARNISH • NO TYPE

ADH
NO COATING

phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur

NO COATING

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ each 15 mL contains: sodium 15 mg ■ store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C red #40, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions?
1-800-719-9260

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Severe Honey Flavor,

Drug Facts (continued)

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than 5 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information

- each 20 mL contains: sodium 14 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, benzyl alcohol, edetate disodium, FD&C red #40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments? 1-800-719-9260

This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Severe Multi-Symptom Cough Cold + Flu.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-16430

100% money back guaranteed.
CVS.com/returnpolicy

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

CVS Health. MAXIMUM STRENGTH

Non-Drowsy
Severe Tussin CF
Adult Cough & Cold

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Compare to the active ingredients in Robitussin® Severe Multi-Symptom Cough Cold + Flu†

CVS Health. MAXIMUM STRENGTH
NDC 51316-323-26

Non-Drowsy
Severe Tussin CF
ACETAMINOPHEN
Pain reliever/Fever reducer
DEXTROMETHORPHAN HBr
Cough suppressant
GUAIFENESIN
Expectorant
PHENYLEPHRINE HCl
Nasal decongestant
Adult Cough & Cold

Relieves:

- Cough, sore throat
- Body aches, fever
- Nasal congestion
- Chest congestion

For Ages 12 Years & Over
Dosage cup provided

Package Contains One Bottle

4 FL OZ (118 mL)

#971824

0 50428 49295 6

28S26 17 C1

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen, USP 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr, USP 20 mg.....Cough suppressant
Guaifenesin, USP 400 mg.....Expectorant
Phenylephrine HCl, USP 10 mg.....Nasal decongestant

Purposes

Uses

- temporarily relieves these symptoms occurring with a cold or flu:
 - cough due to minor throat and bronchial irritation
 - nasal congestion
 - sinus congestion and pressure
 - minor aches and pains
 - sore throat
 - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

CVS Health

Sinus PE Head Congestion + Flu Severe

ACETAMINOPHEN - Pain reliever, Fever reducer
DEXTROMETHORPHAN HBr - Cough suppressant
GUAIFENESIN - Expectorant
PHENYLEPHRINE HCl - Nasal decongestant

Relieves:
 Sinus pressure, Headache,
 Sore throat, Cough, Chest congestion

24 COATED CAPLETS

Actual Size

Compare to the active ingredients in
Sudafed PE® Head Congestion + Flu Severe*

Coated Caplets

Drug Facts (continued)

Directions ■ Do not take more than directed or symptoms.

Keep out of reach of children. In case of overdose, get medical attention or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Warnings ■ If pregnant or breast-feeding, ask a health professional before use. ■ Cough that occurs with too much phlegm (mucus) ■ Difficulty in urination due to enlargement of the prostate gland ■ Chronic bronchitis, or emphysema ■ Persistent or chronic cough such as occurs with smoking, asthma, ■ Cough that occurs with rash or headache that lasts. These ■ Fever gets worse or lasts more than 3 days ■ New symptoms occur ■ Pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ Nervousness, dizziness, or sleepiness occur ■ Stop use and ask a doctor if ■ When using this product do not exceed recommended dosage.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Other Information ■ Each caplet contains: sodium 3 mg

Questions or comments? 1-800-426-9391

Drug Facts (continued)

Warnings ■ Temporarily reduces fever

Do not use ■ With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ heart disease ■ thyroid disease ■ high blood pressure ■ diabetes ■ liver disease

Do not use ■ If you have ever had an allergic reaction to this product or any of its ingredients

Store throat warning: If sore throat is severe, persists for more than 2 days, consult a doctor promptly.

Do not use ■ With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ heart disease ■ thyroid disease ■ high blood pressure ■ diabetes ■ liver disease

Other Information ■ Each caplet contains: sodium 3 mg

Questions or comments? 1-800-426-9391

Drug Facts (continued)

Directions ■ Do not take more than directed or symptoms.

Keep out of reach of children. In case of overdose, get medical attention or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Warnings ■ If pregnant or breast-feeding, ask a health professional before use. ■ Cough that occurs with too much phlegm (mucus) ■ Difficulty in urination due to enlargement of the prostate gland ■ Chronic bronchitis, or emphysema ■ Persistent or chronic cough such as occurs with smoking, asthma, ■ Cough that occurs with rash or headache that lasts. These ■ Fever gets worse or lasts more than 3 days ■ New symptoms occur ■ Pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ Nervousness, dizziness, or sleepiness occur ■ Stop use and ask a doctor if ■ When using this product do not exceed recommended dosage.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Other Information ■ Each caplet contains: sodium 3 mg

Questions or comments? 1-800-426-9391

Drug Facts (continued)

Warnings ■ Temporarily reduces fever

Do not use ■ With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ heart disease ■ thyroid disease ■ high blood pressure ■ diabetes ■ liver disease

Other Information ■ Each caplet contains: sodium 3 mg

Questions or comments? 1-800-426-9391

Drug Facts (continued)

Directions ■ Do not take more than directed or symptoms.

Keep out of reach of children. In case of overdose, get medical attention or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Warnings ■ If pregnant or breast-feeding, ask a health professional before use. ■ Cough that occurs with too much phlegm (mucus) ■ Difficulty in urination due to enlargement of the prostate gland ■ Chronic bronchitis, or emphysema ■ Persistent or chronic cough such as occurs with smoking, asthma, ■ Cough that occurs with rash or headache that lasts. These ■ Fever gets worse or lasts more than 3 days ■ New symptoms occur ■ Pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ Nervousness, dizziness, or sleepiness occur ■ Stop use and ask a doctor if ■ When using this product do not exceed recommended dosage.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Other Information ■ Each caplet contains: sodium 3 mg

Questions or comments? 1-800-426-9391

Drug Facts (continued)

Warnings ■ Temporarily reduces fever

Do not use ■ With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ heart disease ■ thyroid disease ■ high blood pressure ■ diabetes ■ liver disease

Other Information ■ Each caplet contains: sodium 3 mg

Questions or comments? 1-800-426-9391

No Print/No Tamish
Lot and Expiration No.

TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING



#970316

B-0231-546-08-PPCH
REV072254608

PARENTS:
Learn more about our products



100% money back
guaranteed.
CVS.com/returnpolicy

This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Sudafed PE® Head Congestion + Flu Severe.

REV072254608

50844

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2022 CVS/pharmacy
CVS.com® 1-800-SHOP CVS V-19849